

BOLIVIA (PLURINATIONAL STATE OF)

10 725 000

Total population

62%

Percentage of deaths
from NCDs

44 000

Total number of NCD
deaths

16%

Risk of premature death
from target NCDs

1	National NCD targets	●
2	Mortality data	●
3	Risk factor surveys	○
4	National integrated NCD policy/strategy/action plan	○
5	Tobacco demand-reduction measures:	
a	increased excise taxes and prices	○
b	smoke-free policies	●
c	large graphic health warnings/plain packaging	●
d	bans on advertising, promotion and sponsorship	●
e	mass media campaigns	○
6	Harmful use of alcohol reduction measures:	
a	restrictions on physical availability	NR
b	advertising bans or comprehensive restrictions	NR
c	increased excise taxes	NR
7	Unhealthy diet reduction measures:	
a	salt/sodium policies	○
b	saturated fatty acids and trans-fats policies	○
c	marketing to children restrictions	●
d	marketing of breast-milk substitutes restrictions	●
8	Public education and awareness campaign on physical activity	○
9	Guidelines for management of cancer, CVD, diabetes and CRD	○
10	Drug therapy/counselling to prevent heart attacks and strokes	○

● = fully achieved ● = partially achieved ○ = not achieved

NR = No Response

World Health Organization - Noncommunicable Diseases Progress Monitor 2017