Mexico - La Paz (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Mexico-La Paz GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Mexico-La Paz could include in a comprehensive tobacco control program.

The Mexico-La Paz GYTS was a school-based survey of students in 1st, 2nd and 3rd of Secondary and 1st of High School conducted in 2008.

A two-stage cluster sample design was used to produce representative data for Mexico-La Paz. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 88.5%, the class response rate was 98.1%, the student response rate was 79.3%, and the overall response rate was 68.8%. A total of 854 students aged 13-15 participated in the Mexico-La Paz GYTS.

Prevalence

42.7% of students had ever smoked cigarettes (Boys = 40.2%, Girls = 43.0%)

21.9% currently use any tobacco product (Boys = 20.8%, Girls = 20.5%)

13.7% currently smoke cigarettes (Boys = 11.0%, Girls = 14.9%)

11.1% currently use other tobacco products (Boys = 13.2%, Girls = 7.5%)

22.6% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

28.9% think boys and 16.8% think girls who smoke have more friends 12.1% think boys and 10.0% think girls who smoke look more attractive

Access and Availability - Current Smokers

10.9% usually smoke at home 34.9% buy cigarettes in a store

Exposure to Secondhand Smoke (SHS)

38.0% live in homes where others smoke in their presence

50.6% are around others who smoke in places outside their home

87.3% think smoking should be banned from public places

78.2% think smoke from others is harmful to them

42.5% have one or more parents who smoke

13.1% have most or all friends who smoke

Cessation - Current Smokers

49.2% want to stop smoking

60.2% tried to stop smoking during the past year

66.6% have ever received help to stop smoking

Media and Advertising

87.0% saw anti-smoking media messages, in the past 30 days

80.1% saw pro-cigarette ads on billboards, in the past 30 days

71.8% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

16.1% have an object with a cigarette brand logo

11.7% were offered free cigarettes by a tobacco company representative

School

66.8% had been taught in class, during the past year, about the dangers of smoking 36.9% had discussed in class, during the past year, reasons why people their age smoke 62.3% had been taught in class, during the past year, the effects of tobacco use

Highlights

- Over 1 in 5 currently use any form of tobacco; 13.7% of students currently smoke cigarettes; 11.1% currently use some other form of tobacco.
- SHS exposure is high almost 4 in 10 students live in homes where others smoke, and half the students are exposed to smoke around others outside of the home; over 4 in 10 students have at least one parent who smokes.
- More than three-quarters of students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Nearly half of current smokers want to stop smoking.
- One 1 in 6 students has an object with a cigarette brand logo on it.
- Almost 9 in 10 students saw antismoking media messages in the past 30 days; 8 in 10 students saw pro-cigarette ads on billboards and 7 in 10 saw pro-tobacco ads in newspapers or magazines in the past 30 days.