Bolivia - La Paza (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Bolivia - La Paza GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Bolivia - La Paza could include in a comprehensive tobacco control program.

The Bolivia - La Paza GYTS was a school-based survey of students in 1st, 2nd, 3rd, and 4th secondary, and 8th primary conducted in 2000.

A two-stage cluster sample design was used to produce representative data for Bolivia - La Paza. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 95.0%, the class response rate was 100.0%, the student response rate was 83.7%, and the overall response rate was 79.5%. A total of 3,443 students aged 13-15 participated in the Bolivia - La Paza GYTS.

Prevalence

52.4% of students had ever smoked cigarettes (Boy = 59.8%, Girl = 45.6%)

27.3% currently use any tobacco product (Boy = 31.2%, Girl = 23.9%)

23.2% currently smoke cigarettes (Boy = 26.4%, Girl = 20.4%)

9.8% currently use other tobacco products (Boy = 11.7%, Girl = 8.1%)

26.9% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

18.2% think boys and 17.4% think girls who smoke have more friends 13.8% think boys and 12.0% think girls who smoke look more attractive

Access and Availability - Current Smokers

11.0% usually smoke at home

56.7% buy cigarettes in a store

74.7% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

39.8% live in homes where others smoke in their presence

60.6% are around others who smoke in places outside their home

80.9% think smoking should be banned from public places

61.1% think smoke from others is harmful to them

42.6% have one or more parents who smoke

16.3% have most or all friends who smoke

Cessation - Current Smokers

62.6% want to stop smoking

67.7% tried to stop smoking during the past year

67.6% have ever received help to stop smoking

Media and Advertising

75.6% saw anti-smoking media messages, in the past 30 days

88.5% saw pro-cigarette ads on billboards, in the past 30 days

82.0% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

18.5% have an object with a cigarette brand logo

12.5% were offered free cigarettes by a tobacco company representative

School

50.6% had been taught in class, during the past year, about the dangers of smoking

26.7% had discussed in class, during the past year, reasons why people their age smoke

42.2% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 27.3% of students currently use any form of tobacco; 23.2% currently smoke cigarettes; 9.8% currently use some other form of tobacco.
- SHS exposure is high 2 in 5 students live in homes where others smoke, and 3 in 5 students are exposed to smoke around others outside of the home; more than 2 in 5 students have at least one parent who smokes.
- Three in 5 students think smoke from others is harmful to them.
- Eight in 10 students think smoking in public places should be banned.
- Over 3 in 5 current smokers want to stop smoking.
- More than 1 in 5 students have an object with a cigarette brand logo on it.
- Three-quarters of the students saw anti-smoking media messages in the past 30 days; 9 in 10 students saw pro-cigarette ads on billboards and over 4 in 5 saw pro-tobacco ads in newspapers or magazines in the past 30 days.