



**World Health
Organization**



**The WHO STEPwise
approach to stroke
surveillance**

WHO STEPS Stroke Manual

WHO Library Cataloguing-in-Publication Data:

WHO STEPS Stroke Manual: the WHO STEPwise approach to stroke surveillance /
Noncommunicable Diseases and Mental Health, World Health Organization.

1. Cerebrovascular accident - epidemiology. 2. Epidemiologic surveillance - methods. 3. Manuals. I. World Health Organization.

ISBN 92 4 159404 7

(NLM classification: WT 355)

© World Health Organization 2005

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 2476; fax: +41 22 791 4857; email: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press, at the above address (fax: +41 22 791 4806; email: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either express or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Suggested citation:

World Health Organization (2006). WHO STEPS Stroke Manual: The WHO STEPwise approach to *stroke surveillance*. Geneva, World Health Organization.

Acknowledgements:

STEPS Stroke is a collaborative effort between WHO and Non Governmental Organizations (NGOs), notably the International Stroke Society (ISS), the major stroke NGO in official relations with WHO. The Board of the ISS, under the direction of Julian Bogousslavsky, Bo Norrving and Frank Yatsu, has maintained a close interest in the development of surveillance of stroke as part of a broader global stroke initiative. Ruth Bonita chairs the WHO-ISS International Stroke Surveillance Coordinating Unit.

The Stroke surveillance Manual was produced by Ruth Bonita, Peter Heuschmann and Thomas Truelsen. Information Mapping of the manual was undertaken by Charlotte Mill. The Data Entry Tool was developed by the Institute of Epidemiology and Social Medicine, University of Munster, Germany. This WHO Collaborating Centre also provides overall responsibility for data management.

Valuable input and suggestions were provided by the Principal Investigators of the feasibility study undertaken to test the materials in country surveillance sites: Dr Airian, Moscow, Russia; Dr. Dalal, for ICASS centres, India; Dr. Damasceno, Maputo, Mozambique; Dr. Nagaraja, Bangalore, India; Dr. Ogunniyi, Ibadan, Nigeria; Dr. Oveisgharan, Isfahan, Iran; Dr. Pandiyan, Chennai, India; Dr Radhakrishnan Trivandrum, India; Dr. Skvortsava Moscow, Russia. The WHO Regional Advisers and the WHO STEPS Surveillance team are acknowledged for their input into and production of the STEPS surveillance tools (<http://www.who.int/chp/steps>).

Support from the Australian Stroke Society, Canadian Stroke Consortium, European Stroke Conference, Japan Stroke Society, Kenes, Taiwan Stroke Society, the World Federation of Neurology, the World Heart Federation, the World Stroke Federation, and the Government of Finland is also gratefully acknowledged.

The STEPS STROKE Manual can be downloaded at
<http://www.who.int/chp/steps/Stroke/en/>

Table of Contents

SECTION 1: INTRODUCTION

Overview	1-1
Rationale for Stroke Surveillance	1-2
About Stroke	1-4
Major Risk Factors	1-8
In-hospital Management and Facilities	1-9
Overview of STEPS Stroke	1-10
Key Definitions and Concepts	1-12
Setting Objectives	1-13
Stroke Surveillance Process Overview	1-14

SECTION 2: ROLES AND RESPONSIBILITIES

Overview	2-1
Site Coordinator	2-2
Site Data Collection Staff	2-3
International Coordination and Support	2-4

SECTION 3: PLANNING AND PREPARING A STROKE STUDY

Overview	3-1
Process Overview and Eligibility	3-2
Identifying the Scope	3-3
Defining the STEPS Stroke Surveillance Site	3-5
Identifying the Register Population	3-7
Modifying the Stroke Instrument	3-10
Registration for participation in STEPS Stroke	3-11
Getting Ethical Approval	3-13

SECTION 4: PREPARING THE STROKE SURVEILLANCE SITE

Overview	4-1
Recruiting Staff	4-2
Briefing and Training for Data Collection Staff	4-3
Setting up the Stroke Surveillance Site	4-4
Installing and Preparing the Data Entry Tool	4-5
Test Run	4-7

Continued on next page

Table of Contents, Continued

SECTION 5: DATA COLLECTION GUIDELINES

Overview	5-1
Case Finding Methods.....	5-2
Identifying Stroke Patients in Hospitals (Step 1)	5-3
Identifying Fatal Stroke Patients in the Community (Step 2)	5-5
Estimating Non-Fatal Stroke Events in the Community (Step 3)	5-7
Interview Skills	5-9
Recording Responses for Registration	5-11
Completing the Stroke Instrument	5-12
Guide to Completing All Stroke Events.....	5-13
Guide to Completing Step 1: Events Admitted to Hospital	5-15
Guide to Completing Step 2: Fatal Community Events.....	5-22
Guide to Completing Step 3: Non-Fatal Community Events.....	5-22

SECTION 6: DATA ENTRY AND DATA MANAGEMENT

Overview	6-1
Data Entry	6-2
Data Management	6-5
Creating Reports.....	6-7
Exporting Data	6-8

SECTION 7: STEPS STROKE INSTRUMENT AND FORMS

STEPS Stroke Instrument.....	7a-1
Registration for Participation	7b-1

SECTION 8: GLOSSARY AND REFERENCE MATERIAL

Glossary of Terms Used in STEPS Stroke.....	8-2
Source Publications and References	8-3

Section 1: Introduction

Overview

Introduction This section is an introduction to the STEPS Stroke Surveillance Manual.

Purpose The purpose of the manual is to provide guidelines and supporting material for sites embarking on STEPS stroke surveillance, so they are able to:

- plan and prepare the surveillance register scope and environment
 - recruit and train data collection staff
 - establish and maintain the stroke register
 - report and disseminate the results.
-

Intended audience The manual is primarily intended for the stroke surveillance site principal investigator. Parts of the manual are also intended for data collection staff.

Guide to using the manual The manual has been written in modular parts and is structured to follow the sequence of events required to implement a STEPS Stroke register. It is divided into eight sections. Each section is introduced with a table of contents to help readers find specific topics. The manual includes both general information and specific instructional material that can be extracted and used for:

- training
- data collection
- data entry

Page numbers have two components. The first number refers to the section and the second to the page number in that section.
For example: 3-6 indicates section 3, page 6.

In this section This section contains the following topics:

Topic	See Page
Rationale for Stroke Surveillance	1-2
About Stroke	1-4
Major Risk Factors	1-8
In-hospital Management and Facilities	1-9
Overview of STEPS Stroke	1-10
Key Definitions and Epidemiological concepts	1-12
Setting Objectives	1-13
Stroke Surveillance Process Overview	1-14

Rationale for Stroke Surveillance

Introduction	Well-conducted stroke surveillance (with accurate and complete registers) provides essential data that can be used to improve appropriate allocation of health resources.
Definition of surveillance	<p>Surveillance is the ongoing, systematic:</p> <ul style="list-style-type: none">• collection• analysis• interpretation, and• dissemination of health information.
Purpose of STEPS stroke surveillance	<p>The purpose of the STEPS stroke surveillance register is to provide health workers and policy makers with a standardized tool to:</p> <ul style="list-style-type: none">• assess the magnitude of stroke• describe populations at risk• identify associated risk factors• monitor trends over time• provide the basis for designing and implementing interventions• monitor and evaluate the effectiveness of interventions.
The evidence	<p>Globally, cerebrovascular disease (stroke) is the second leading cause of death. It is a disease that predominantly occurs in mid-age and older adults.</p> <p>WHO estimated that in 2005, stroke accounted for 5.7 million deaths world wide, equivalent to 9.9 % of all deaths. Over 85% of these deaths will have occurred in people living in low and middle income countries and one third will be in people aged less than 70 years.</p>
Rationale for surveillance	<p>While many countries struggle with the consequences and problems of communicable diseases, chronic noncommunicable diseases are on the rise. In addition to being a major cause of death, many surviving stroke patients are disabled and need help in activities of daily living, which must be provided by family members, the health system, or other social institutions.</p> <p>Lack of data on stroke from many countries hampers efficient coordination of stroke prevention, treatment, and rehabilitation. Due to ongoing demographic changes, strategies to reduce future stroke burden and ensure adequate health resources are urgently needed. STEPS stroke surveillance provides the framework for data collection and comparisons between and within populations.</p>

Continued on next page

Rationale for Stroke Surveillance, Continued

Other expected outcomes

Other expected outcomes of setting up surveillance sites include:

- Building expertise and high quality systems for long-term community surveillance of chronic noncommunicable diseases, especially stroke
 - Establish a research network
 - Increase awareness of noncommunicable disease in the community.
 - Establish country-specific priorities for the prevention and management of stroke in the context of national integrated plans for chronic disease prevention and control (see http://www.who.int/chp/chronic_disease_report)
-

Surveillance: key to prevention

Above all, clinical trials and epidemiological studies have shown that stroke to a large extent is preventable. Public actions to lower the prevalence of exposure to risk factors, however, are unlikely to be taken, if the magnitude and consequences of stroke and other major chronic diseases are not identified.

Prevention strategies

Once reliable data are available, different prevention strategies can be implemented to reduce the occurrence and impact of stroke as described in the table below:

Prevention strategy	Aimed at reducing..	For example, through..
Primary	Occurrence of stroke in the first place.	<ul style="list-style-type: none">• Identification of individuals at high overall risk of stroke or CVD (hypertensive people or diabetics)• Population wide initiatives to increase physical activity• Legislation to control tobacco use.
Secondary	Impact of stroke in people who already suffer from a stroke or TIA.	<ul style="list-style-type: none">• Intensified reduction in exposure to major cardiovascular risk factors• Anti-platelets and anti-hypertensive treatment.
Tertiary	Consequences and damages in stroke patients.	<ul style="list-style-type: none">• Treatment of infections in the acute stage• Management of co-morbidities• Improved rehabilitation.

About Stroke

Introduction

Stroke is a clinically defined disease making it possible to capture data and follow trends in incidence or hospital admission rates in many different countries irrespective of access to technological equipment.

A costly disease

Stroke is a costly disease because of the:

- Large numbers of premature deaths,
 - Ongoing disability in many survivors,
 - Impact on families or caregivers and
 - Impact on health services.
-

Standard WHO definition

The recommended standard WHO stroke definition is as follows:

A focal (or at times global) neurological impairment of sudden onset, and lasting more than 24 hours (or leading to death), and of presumed vascular origin.

This clinical definition has four components:

- A neurological impairment or deficit of
 - Sudden onset, and
 - Lasting more than 24 hours (or leading to death), and
 - Of presumed vascular origin.
-

Notes on the standard definition

The WHO standard definition excludes:

- Transient ischemic attack (TIA), which is defined as focal neurological symptoms but lasting less than 24 hours
- Subdural haemorrhage
- Epidural haemorrhage
- poisoning
- Symptoms caused by trauma.

“Global” refers to patients with subarachnoid haemorrhage or deep coma but excludes coma of systemic vascular origin such as:

- Shock
- Stokes-adams syndrome
- Hypertensive encephalopathia.

Stroke is a clinical diagnosis and not based on radiological findings

Continued on next page

About Stroke, Continued

Types of stroke There are three major stroke sub groups as follows:

- Ischemic stroke
- Intracerebral haemorrhage
- Subarachnoid haemorrhage.

Type	Caused by	Diagnosis based on
Ischemic stroke	Sudden occlusion of arteries supplying the brain. Due either to a thrombus formed: <ul style="list-style-type: none">• Directly at the site of occlusion (thrombotic ischemic stroke), or• In another part of the circulation, which follows the blood stream until it obstructs arteries in the brain (embolic ischemic stroke).	<ul style="list-style-type: none">• Neuro imaging recordings Note: it may not be possible to decide clinically or radiological whether it is a thrombotic or embolic ischemic stroke.
Intracerebral haemorrhage	Bleeding from one of the brain's arteries into the brain tissue. Note: May be more prevalent in developing countries possibly due to diet, physical activity, insufficient treatment of raised blood pressure, and genetic predisposition.	<ul style="list-style-type: none">• Neuro imaging recordings
Subarachnoid haemorrhage	Arterial bleeding in the space between the two meninges, pia mater and arachnoidea. Note: Typical symptoms are sudden onset of very severe headache and usually impaired consciousness.	<ul style="list-style-type: none">• Neuro imaging, or• Lumbar puncture.

Note: Each type differs with respect to survival and long-term disability.

General major symptoms Symptoms should be of a presumed vascular origin and should include one or more of the following definite focal or global disturbances of the cerebral function:

- Unilateral motor impairment (including lack of coordination)
- Unilateral sensory impairment
- Aphasia/dysphasia (non-fluent speech)
- Hemianopia (half-sided impairment of visual fields)
- Forced gaze (conjugate deviation)
- Apraxia of acute onset
- Ataxia of acute onset
- Perception deficit of acute onset.

Continued on next page

About Stroke, Continued

Other symptoms

Other symptoms that may be present but are not adequate for stroke diagnosis (often resulting from other diseases or abnormalities such as dehydration, cardiac failure, infections, dementia, and malnutrition) are as follows:

- Dizziness, vertigo
 - Localized headache
 - Blurred vision of both eyes
 - Diplopia
 - Dysarthria (slurred speech)
 - Impaired cognitive function (including confusion)
 - Impaired consciousness
 - Seizures
 - Dysphagia.
-

Subarachnoid haemorrhage

For subarachnoid haemorrhage at least one of the following must be present in addition to the general major symptoms:

- Recent subarachnoid hemorrhage, aneurysm or arteriovenous malformation (necropsy/autopsy).
 - Blood in the Fissura Sylvii or between the frontal lobes or in the basal cistern or in cerebral ventricles (CT or MRI).
 - Blood stained cerebrospinal fluid ($>2\,000$ red blood cells per mm^3), aneurysm or an arteriovenous malformation (angiography).
 - Blood stained cerebrospinal fluid ($>2\,000$ red blood cells per mm^3), also xanthochromic and intra-cerebral haemorrhage (necropsy or CT).
-

Stroke like symptoms

A broad range of other diseases may cause similar symptoms, for example:

- HIV/AIDS
- Tuberculosis
- Syphilis
- Intracerebral cancer.

These diseases are known to be able to cause focal neurologic disturbances and thereby mimic a stroke. Attention to the development of symptoms is an important factor to consider in order to avoid other diseases being misinterpreted as vascular disease, and leading to ineffective preventive strategies.

Continued on next page

About Stroke, Continued

Types of stroke events

There are four types of stroke events:

Type of stroke event	Defined as occurring in a person who has..
First ever	First ever (also called "first in a lifetime") refers to people who have never had a stroke before. Note: Previous TIA is not considered a stroke as symptoms last less than 24 hours.
Recurrent	There are two types of recurrent strokes: <ul style="list-style-type: none">• A history of a previous stroke event at some time in the past which meets the WHO definition, and• A history of a new stroke event occurring more than 28 days after onset of a stroke event already registered.
Non-fatal	A stroke case who survived at least 28 days after the onset of the stroke symptoms.
Fatal	Died within 28 days of stroke symptom onset.

First ever stroke events

A first ever stroke event is the same as first in a lifetime event. (see Key definitions and epidemiological concepts Section 1-12)

Recurrent stroke events

For a new episode of symptoms to be counted as a new or recurrent stroke event, general stroke criteria as defined above must be met and either:

- The previous event in the same arterial distribution must have occurred 29, or more days previously (by subtraction of dates), or
- The new event is unequivocally in a different arterial territory than an earlier event which has occurred 28 or fewer days previously.

If a patient experiences further acute symptoms suggestive of stroke within 28 days of the onset of a first episode and in the same carotid or vertebral artery territory, this second episode is **not** counted as a new stroke event.

Equally, if a patient experiences further acute symptoms suggestive of stroke after 28 days of the onset of a first episode, this second episode **is** counted as a new stroke event.

Note: Each **event** is registered separately. This means that 2 (or more) forms will be completed for the same individual who experiences multiple new acute stroke events that meet the criteria.

Major Risk Factors

Introduction Stroke is a multi factorial disease where a combination of risk factors, all of which do not all have to be present, will, over time, influence the subject's likelihood of suffering a stroke.

Major risk factors The major risk factors can be divided into the following categories:

Category	Risk factors
Modifiable	<ul style="list-style-type: none">• Elevated blood pressure• Tobacco use• Physical inactivity• Diet (low fruit and vegetable consumption)• Heavy alcohol consumption• Overweight• Diabetes
Environmental	<ul style="list-style-type: none">• Passive smoking• Access to medical treatment.
Non-modifiable	<ul style="list-style-type: none">• Age• Sex (eg. high age and male sex are in many populations associated with an increased risk).• Family history; genetics

Other risk factors In developed countries, diabetes mellitus as well as atrial fibrillation and other cardiac diseases are other important modifiable risk factors for ischemic stroke.

The role of hypercholesteremia as risk factor for stroke is currently part of an ongoing discussion. There is evidence that lower total cholesterol levels might be associated with a decreased risk of ischemic stroke but also might be accompanied by higher rates of hemorrhagic strokes.

Key sources on information on country level risk factors WHO has developed two major tools for collecting, displaying and analysing data on the major 8 major risk factors which are common to CVD.

- The STEP wise framework for surveillance of Risk Factors for chronic diseases; see website: <http://www.who.int/chp/steps>
- The Global NCD InfoBase provides biennial updates on available country-level risk factor data including comparable mortality estimates for stroke.

See website: http://www.who.int/ncd_surveillance/infobase

In-hospital Management and Facilities

Introduction In-hospital management refers to stroke patients admitted to a health facility.

Specialist stroke teams Stroke patients admitted to a hospital department with a specialized stroke team or multi-disciplinary approach have a better outcome than patients admitted to departments without such teams or approach. This is measured in terms of long term reduction of death, and of dependency and institutionalisation.

Early rehabilitation and early mobilization of patients with severe neurological deficits helps lower disability after stroke and avoids complications.

Medication Different treatments and medications identified in the STEPS Stroke instrument have been shown to reduce risk of stroke in selected groups of patients in predominantly high-income countries. These are explained in the following table.

Type of medication	Used to
New and old anti-hypertensive drugs	Lower blood pressure and reduce stroke occurrence.
Aspirin, Dipyridamole, and Clopidogrel	Reduces platelet aggregation and reduces risk of a new ischemic stroke.
Anti-coagulant therapy	Prevent cardiac embolism in patients with atrial fibrillation.
Intravenous treatment with tissue plasminogen activator (tPA)	Dissolves the blood clots in patients with acute ischemic stroke.

Overview of STEPS Stroke

Introduction

The STEPwise approach to stroke surveillance provides a flexible system and an opportunity for all countries to get started and contribute and share data on stroke.

Basis of STEPS Stroke

STEPS Stroke identifies three different groups of stroke patients who make up the burden of stroke in any given community or population. They are listed in the order of complexity of identifying them:

- Information on stroke patients admitted to health facilities (Step 1);
- Identification of fatal stroke events in the same community (Step 2);
- Estimates of non-fatal stroke events in the same community (Step 3).


Within each Step (1, 2 and 3) there are a further two possible levels of information that can be collected (Core, and Expanded).

By using the same standardized approach, all countries can monitor trends within countries and between countries.

The STEPS Stroke instrument was developed, in part, by using the protocol from the WHO MONICA Project.

STEPS Stroke diagram

The following diagram illustrates the general concept of the WHO STEPwise approach



Note: An "ideal" stroke incidence register requires that all eligible stroke occurring are identified in residents of the defined source population,

Continued on next page

Overview of STEPS Stroke, Continued

STEPS stroke tools

A suite of tools have been developed to help methodically and consistently work through the STEPS surveillance process. This suite of tools is called the STEPS Stroke Starter Kit and includes a:

- STEPS Stroke manual
 - STEPS Stroke instrument
 - Forms and templates
 - Data entry tool (to create a stroke register).
-

STEPS stroke instrument

The STEPS stroke instrument is a standardized questionnaire used to collect stroke patient data to be entered into the register using the data entry tool.

The STEPS stroke instrument covers three different 'Steps' of stroke case finding (Step 1, Step 2 and Step 3) for a defined population. Within each Step, there are three different levels of data collection of increasing complexity (Core, Expanded, Optional) as follows:

Step		Core	Expanded	<i>Provides data on</i>
1	Hospitalized events (fatal and non-fatal)	<ul style="list-style-type: none">• Demographic information• Time of onset• Vital status day 10	<ul style="list-style-type: none">• Treatment• Disability• Type of stroke	<i>Stroke admissions and hospital case fatality</i>
2	Fatal events in the community	<ul style="list-style-type: none">• Demographic information• Death certificates, or• Verbal autopsy	<ul style="list-style-type: none">• Autopsy/necropsy reports• Type of stroke	<i>Stroke deaths</i>
3	Non-fatal events in the community	<ul style="list-style-type: none">• Demographic information• Time of onset• Vital status day 10	<ul style="list-style-type: none">• Treatment• Disability• Type of stroke	<i>Stroke incidence and case fatality</i>

Recommended steps

The optimal stroke surveillance system requires collection of data from all three steps and provision of census data from the source population.

Costs and complexity increase with identifying subgroups of patients at each step. The level of complexity will therefore depend on development of health services and resources, and each participating country may collect the amount of data that it finds is feasible. Not all sites will be able to undertake an "ideal" incidence register.

Continued on next page

Some Key Definitions and Epidemiological Concepts

Prevalence and Incidence

Prevalence and incidence are fundamentally different ways of measuring the occurrence of a disease although both involve the counting of cases in defined populations at risk. The defined population includes people living within a geographically defined area that a particular hospital or hospitals serve. The defined population, depending on the level of analysis, can be further observed by additional characteristics (age, sex, ethnicity, nationality, residency, etc.)

- The prevalence of stroke is the number of cases in a defined population at a specified point in time - gives a "snapshot" of survivors at any one time
- The incidence is the number of new cases of stroke arising in a given period in a defined population - gives an indication of the risk of stroke.

Data on both prevalence and incidence become much more useful if converted into rates. A rate is calculated by dividing the number of cases by the corresponding number of people in the defined population at risk.

Calculating the Incidence rate

Incidence rate (*I*) is calculated as follows:

$$I = \frac{\text{Number of people who get a stroke in a specified time}}{\text{Number of people in the population at risk over the length of time during which each person in the population is at risk}} \times (10)^5$$

Case series

A case series refers to stroke cases identified in specified hospital facilities (numerator) but without reference to a defined population from which they came (denominator). Without a denominator, rates can not be calculated. Even so case series present important clinical information about stroke, perhaps for the first time. Case series are often the beginning of the process which helps establish more robust measures subsequently.

Surveillance: ongoing

While a stroke register can be a one off exercise, surveillance involves commitment to developing the stroke register on an ongoing and /or repeated basis. This may also be in the form of repeat studies (every 5 to 10 years), particularly to look at hospital or population trends.

It is recommended that when a STEPS Stroke Surveillance register is launched for the first time, there should be a plan for future follow-up to measure trends. This can be achieved by continuous surveillance as part of a broader health information system

Setting Objectives

Long term monitoring of stroke

Establishing a surveillance system for long term monitoring of stroke can help to:

- Understand the clinical profile of stroke patients
 - Develop complete, population-based systems of case ascertainment for stroke
 - Verify the accuracy of routine data systems for monitoring stroke
 - Develop local expertise in epidemiological research methods through education and training.
-

Size and impact of the problem

Undertaking stroke surveillance also helps to determine the size of the problem and help to determine the following:

- Incidence, case fatality and outcome of stroke
 - Impact on health care systems of stroke
 - Use of acute hospital beds, rehabilitation facilities, and requirements for community care including impact on families of stroke
 - Use of effective interventions in the acute phase (stroke unit, use of drugs etc).
-

Stroke Surveillance Process Overview


Introduction

For STEPS Stroke Surveillance to be effective, the whole process needs to be properly planned and organized before being implemented. Guidelines are provided below to help you plan a STEPS stroke surveillance register.

Key stages, tasks and timeframes

The minimum recommended total timeframe to collect data for a STEPS stroke register is 12 months.

The chart below shows each of the main stages and tasks in a STEPS Stroke register with indicative timeframes for each phase and task.



Section 2: Roles and Responsibilities


Overview

Introduction There are a number of entities involved in coordinating and implementing STEPS Stroke surveillance. Representation is covered at the:

- country
 - sub-regional, and
 - regional level.
-

Purpose The purpose of this section is to provide an overview of the core roles involved in STEPS Stroke surveillance.

STEPS stroke surveillance network The diagram below shows how the global Stroke surveillance network is organised.



In this section This section covers the responsibilities for the following roles:

Topic	See Page
Site Coordinator	2-2
Site Data Collection Staff	2-3
International Coordination and Support (ICS)	2-4

Site Coordinator

Introduction

The STEPS stroke surveillance (SSS) site coordinator is the local principal investigator. This key person is responsible for planning and coordinating the local STEPS Stroke surveillance register.

The site coordinator should be familiar with the entire manual to understand the whole STEPS stroke surveillance process.

Skills and attributes

The site coordinator will need to have the following qualifications and, general skills and attributes:

- Neurological or stroke physician (or nurse) with proven experience in the field of cerebrovascular disease.
 - Good understanding of epidemiological principles of differences between hospital based stroke registers and population based stroke registers
 - Good understanding of the general philosophy and objectives of the global STEPS Stroke surveillance process.
 - Good written and oral communication skills and proficient in English.
 - Ability to recruit and train interviewer staff.
-

Level of authority

The site coordinator should have sufficient authority to:

- Negotiate and obtain resources for the whole stroke register.
 - Oversee progress of the national, sub-national, district or local STEPS stroke surveillance implementation.
 - Contribute to the disease prevention and health promotion activities that will arise from the data gathered by STEPS Stroke.
-

Core roles

The core roles of the site coordinator may include all or some of the following:

Role	Description
1	Planning and preparing for a STEPS Stroke register.
2	Applying for participation.
3	Identifying and securing local funding and / or "in kind" support.
4	Handling ethical approval.
5	Recruiting and training interview staff.
6	Supervising data collection and adjudicating difficult diagnoses.
7	Reporting results.
8	Planning and preparing for future studies.
9	Liaising with the International coordination and support (ICS), local authorities, WHO/PAHO NCD regional, country and CAREC representatives and other stakeholders.
10	Completing test stroke cases provided by ICS for quality control purposes.

Site Data Collection Staff

Introduction The STEPS Stroke surveillance (SSS) site data collection staff are all those who have been trained to collect the data and enter it into the stroke register.

Interviewer roles Data collection roles will depend on the scope of the register. Core roles for a data collection staff member may include all or some of the following. Specific tasks are identified in section 5.

Role	Description
1	Actively identifying stroke patients admitted to (or occurring within) the hospital on a daily or weekly basis.
2	Retrospectively reviewing records of stroke patients.
3	Resolving difficult cases (where a patient needs to be assessed by an experienced medical practitioner or neurologist).
4	Recording patient details on the Stroke instrument
	Entering instrument data into the register (using the data entry tool)
5	Following up with patients at day 28.
6	Liaising with and reporting any difficulties to the site coordinator.

Skills and attributes Interviewers should have the following general skills and attributes:

- Good basic knowledge of different clinical symptoms of stroke.
 - Good understanding of the different case finding methods (hot and cold pursuit).
 - Excellent understanding of the stroke definition and the instrument questions.
 - Have a sensitive approach towards people who are in a stressed situation or are recalling a sad moment in life.
 - Good oral, written and keyboard skills.
 - Good attention to detail.
 - Ability to follow instructions consistently but raise concerns when appropriate.
 - Work well with others to achieve results.
-

International Coordination and Support (ICS)

Introduction	PAHO/WHO and CAREC provide technical support and guidance for STEPS stroke surveillance in the Americas and the Caribbean.
Objectives	The main objective is to oversee the practical and logistic issues relating to the overall coordination and implementation of STEPS Stroke surveillance.
Core roles	<p>The core roles of PAHO and CAREC are to:</p> <ul style="list-style-type: none">• register surveillance site participation• support the site coordinator• provide access and support to STEPS surveillance tools and reference material• share updates periodically• oversee the overall implementation of the STEPwise approach to stroke surveillance (STEPS Stroke)• ensure quality control• provide feedback on all drafts of reports before they are submitted.
Support	<p>PAHO Regional program has supported dissemination of the Methodology in the Region, through translation of the Manual in Spanish and Portuguese (http://www.paho.org/English/AD/DPC/NC/steps-stroke.htm) and promoting at international meetings. PAHO and CAREC offer temporary advisors for consultancy. PAHO and CAREC are responsible for advocacy around STEPS Stroke and overseeing the practical and logistic issues relating to the overall implementation of STEPS Stroke. The core responsibilities are to:</p> <ul style="list-style-type: none">• Advocate at major international stroke meetings.• Develop closer links between the major NGOs, PAHO/WHO and CAREC around stroke surveillance.• Help expand the number of Stroke surveillance sites.• Liaise with site coordinators on a regular basis.• Support and maintain the linkages on the STEPS web page.• Receive requests for participation and refer to the ICS technical adviser.• Support participating stroke surveillance sites (SSS) with general information about the manual.• Provide technical support to participating SSS.• Ensure relevant PAHO/WHO regional and country people are informed.• Update and maintain the STEPS Stroke surveillance manual.• Develop and maintaining Stroke Data Entry Tool (DET) in collaboration with the temporary adviser.• Modify the Stroke DET in accordance with experience from the feasibility register of the SSS.• Collect and collate data for the annual report and for conference presentations as necessary.

Section 3: Planning and Preparing a Stroke Register

Overview

Introduction This section covers the tasks that need to be conducted to plan and prepare for STEPS stroke surveillance.

Intended audience This section is primarily designed to be used by those fulfilling the role of the Site coordinator and associated advisory group.

Using existing case registration systems In some settings, there will be other hospital-based chronic disease case registration systems that cover large populations. Where these systems already exist, consider working with the registration teams and adding stroke surveillance to their work.

In this section This section covers the following topics:

Topic	See Page
Process Overview and Eligibility	3-2
Identifying the Scope	3-3
Defining the STEPS Stroke Surveillance Site	3-5
Identifying the Register Population	3-7
Modifying the Stroke Instrument	3-9
Registering for Participation	3-11
Getting Ethical Approval	3-13

Process Overview and Eligibility

Introduction

Before registering for participation in STEPS Stroke (see section 7b), some initial prerequisite actions and criteria must be defined.

Process overview

The table below shows each stage in the planning, scoping and eligibility process.

Stage	Description						
1	Define the type and scope of the register (Step 1, 2, 3). The three options are as follows: <ul style="list-style-type: none">- a hospital register case series (Step 1 with no population base)- a hospital register linked to a defined population base (Step1)- an incidence register linked to a defined population base (Steps 1-3)						
2	Identify the register site.						
3	Identify the defined population from which the cases will be derived (see Section 1-12) <table><tr><th>If the source population is</th><th>Then</th></tr><tr><td>Available</td><td>Apply for full participation</td></tr><tr><td>Not available</td><td>Apply for limited participation (case series only)</td></tr></table>	If the source population is	Then	Available	Apply for full participation	Not available	Apply for limited participation (case series only)
If the source population is	Then						
Available	Apply for full participation						
Not available	Apply for limited participation (case series only)						
4	Prepare the instrument.						
5	Obtain sustainable funding.						
6	Apply for participation.						
7	Get ethical approval.						

Note: Each of these stages is explained in more detail below.

Identifying the Scope

Introduction

The focus of STEPS Stroke surveillance is reflected in the core of the stroke instrument. All countries should be able to undertake the core items of Step 1, although not all countries will have access to the defined population from which the stroke events arise.

Stroke register design

The table below provides an overview of the different designs of a STEPS Stroke register. The usefulness of the register is influenced by the quality, completeness and population coverage. A case series poses the greatest challenge in interpretation, but may be the only option in those countries where there are no census data for the catchment area covered by the selected hospitals. (see Section p1-12, 1-13).

Defined Population available	↑	Step 3		Incidence rates (hospital + fatal& non fatal strokes in the community)		
		Step 2			Case fatality (hospital events + fatal strokes in the community)	
		Step 1				Hospital admission rates
No Defined Population Available		Step 1				Case series

Step 1 data collection

Step 1 focuses on residents (preferably of a defined population) who are admitted to a health facility (hospitals, nursing homes or other similar health facilities) with a stroke which meets the WHO definition. A hospital based stroke register provides data on:

Step 1	Register of
Core	<ul style="list-style-type: none"> Stroke admissions Severity of stroke Survival rates for this group of patients
Expanded	<ul style="list-style-type: none"> Pre-stroke exposure to major risk factors

Step 1: Main outcomes

The main outcomes from this Step include:

- Health facility resources allocated to stroke patients
- Functional status of stroke patients at discharge.
- Risk factor exposure.
- Hospital admission rates when combined with population estimates from which the stroke patients are derived.

Note: Step 1 alone does not provide estimates of stroke incidence in the population because some patients die before hospital admission can be arranged and others are cared for in the community rather than in hospital.

Continued on next page

Identifying the Scope, Continued

Step 2: Data collection

Step 2 builds on the hospital register in Step 1 by validating death certificates from routine sources to include fatal stroke events that have occurred in the same community but out of hospital. These data are derived from death certificates need to be validated by verbal autopsy (see section 5-5, 5-6).

Step 2: outcomes

The main outcome from Step 2 (combined with the data from the hospital register from Step 1) is calculation of specific mortality rates and years of life lost due to stroke in the defined population. These can be broken down by:

- Age and sex
 - Proportion of fatal events occurring outside of health facilities
 - Years of life lost because of stroke (YLLs).
-

Step 3: Data collection

The surveillance of stroke is complicated by the fact that a high number of cases are not admitted to hospital. Step 3 is therefore the most challenging subset of eligible patients to identify. Their identification is vital for the accurate determination of stroke incidence. These strokes are a combination of milder and more severe strokes than those that come to hospital, and consequently their inclusion influences case fatality.

Step 3: outcomes

The main outcome from Step 3 (combined with Step 1 and Step 2 results), is the calculation of incidence and case-fatality. It also allows estimates of:

- Stroke incidence, prevalence, and case fatality
 - Years of Life lived with Disability (YLDs)
 - Estimate of needs for long term care.
-

Recommended scope

The minimum recommended scope for most countries should be Step 1 - preferably with a well defined source population from which eligible patients are derived.

Some countries will be able to achieve a population based register (involving all 3 steps). This provides the most valuable epidemiological measures for public health initiatives for stroke prevention. It is therefore recommended that there is an intention to advance the register to include all three Steps or subgroups of patients, if resources allow and access to central death certificates is available.

Financial support

Once you have identified the scope of your register, you will need to set out a budget and seek financial support (from local or national sources or in kind) to cover all expenses of the register for the whole register period.

Defining the STEPS Stroke Surveillance Site

Introduction	<p>The next stage in the process of being eligible to participate includes identifying and/ or describing the STEPS stroke surveillance site. This may differ depending on the type of register being planned: case series or one which produces hospital rates.</p>
Step 1: hospital based register	<p>When developing a hospital based register to be linked to a population, all health facilities, or network of health facilities that are found within the defined population need to be identified and involved in the register. These could include:</p> <ul style="list-style-type: none">• All health facilities in the (source population) area• A small group of health facilities that admit most stroke events• Wards within defined health facilities that admit most stroke events. <p>Note: To define the health facilities, complete the hospital information form in section 7b . Once defined, the (group of) selected health care facilities, together with the source population, will be referred to as the STEPS stroke surveillance site (SSS).</p>
Step 2: strokes dying outside hospital	<p>Key case finding sources include access to routine death certificates and an ability to verify all deaths possibly due to a stroke event (including "old age") by use of verification using verbal autopsy techniques.</p>
Step 3: events cared for only in the community	<p>Key case finding sources for stroke cases cared for entirely at home, involves the collaboration and cooperation of, among others, to ensure ongoing support and referral of eligible cases to the register and may involve the following:</p> <ul style="list-style-type: none">• General practitioners and other health care providers in the community who need to notify the register team of all such events• Community health nurses and village elders/church leaders or• Alternative medicine practitioners, faith healers etc.
Defining the population	<p>Once you have defined the source population or community (preferably from the most recent census) from which the stroke cases will be identified, send the ICS a copy with the registration form (see section 7b).</p>
Patient eligibility	<p>A patient is eligible for inclusion in the stroke register, if:</p> <ul style="list-style-type: none">• A resident in the defined population of the stroke surveillance site,• Meets the age selected (see Section page 3-8)• Has a stroke event within the defined period of time

Continued on next page

Defining the STEPS Stroke Surveillance Site, Continued

Estimation of expected stroke events

To be eligible to participate, a minimum of 250 stroke patients per year in the source population from which the cases will be derived, (i.e. hospital and / or community) is necessary in order to

- Ensure meaningful analysis of the data by age and sex
- Have sufficient numbers to detect trends over time.

These can be based on previous experience or the results of a pilot study.

Identifying the Register Population

Introduction	<p>Calculation of epidemiologic rates is based on the number of stroke events occurring in the defined population at risk. The ideal population comes from a well defined geographic area.</p> <p>One of the first steps in setting up surveillance registers is therefore to specify and describe the population in which the register is going to take place.</p>
Requirement	<p>A defined source population should include population counts broken down by:</p> <ul style="list-style-type: none">• Each age group to be included in the stroke surveillance register• Sex, and• Total counts.
Source of information	<p>In many settings, source population counts can be obtained from:</p> <ul style="list-style-type: none">• Population census lists• Inter census estimates• Population registers.
Where source population data does not exist	<p>In settings where data for a well-defined geographic population does not exist, you will only be able to produce a case series stroke register.</p> <p>Interpretation of this data over time poses major challenges because of changing hospital practices and because of lack of information about the nature of the population from which the cases come.</p>
Balancing population coverage and number of stroke cases	<p>Sites that wish to estimate admission rates for Step 1 and/or do Step 2 and Step 3 must provide an accurate estimate of the defined population at the time of registration.</p> <p>To provide a reliable estimate of the impact of stroke occurrence, representative regional population coverage (from around 250,000 total population up to 1 million) is recommended.</p> <p>Including more than 1 million people is usually not possible and would require a sample system to be established and a much larger team than the one recommended in this Manual.</p>

Continued on next page

Identifying the Register Population, Continued

Factors to consider

The table below lists some factors to consider for population coverage.

Coverage	Guidelines
Districts	Consider both urban and rural. Note: Often there are differences between urban and rural districts with respect to exposure to risk factors, treatment of predisposing diseases, for example hypertension, and access to health authorities and facilities.
Hospitals	Include both private and public (state run).
Gender	Include both men and women.
Socioeconomic status	Allow a representative range of socioeconomic groups.

Age range

For practical and financial reasons, you should restrict the Core register to age groups where stroke usually occurs, for example from age 30 to 84 years.

If you need to expand the register to assess stroke cases in the very young or very old, you may wish to include other age ranges. See the table below for guidance on expanded and optional age ranges.

STEPS levels	Age range (years)
Core	35-84
Expanded	15 - 34 85 +

Note: It is often difficult to determine actual stroke in the very elderly due to co-morbidity. Including the expanded age range 85+ can therefore skew results.

Sex

Stroke rates are often higher in men than in women, although the differences are not as marked as for other chronic diseases (such as heart disease). Men and women should be presented separately in all analyses.

Describing the register population

For a stroke surveillance site intending to undertake complete coverage of possible stroke events occurring in residents in the defined population, details are requested as part of the process of registering for participation in STEPS Stroke Surveillance. This form is available in section 7b.

Modifying the Stroke Instrument

Introduction The Stroke Instrument is a standard document that allows comparisons and international trends analysis and should not be changed. It uses a standard international calendar and is the basis for the standard data entry tool.

Minor modifications Despite the need for standardization, some minor local modifications may be required in some settings (for example, to clarify terminology, or provide a more comprehensive assessment of stroke occurrence and treatment). The following table provides guidance on possible situations where the Instrument may be modified to local requirements.

Modification type	If...	Then...
Terminology	The terms used in some Core 'standard' questions do not fit the cultural setting (for example, ethnicity).	Alter the term for local relevance, but ensure the original meaning is retained.
Additional information	You require additional data on stroke occurrence and treatment (for example, use of tPA) and you have available resources.	Add selective, but limited questions as Optional items.
Link to previous data	You require specific data to link to previous surveys	Add selective, but limited questions as Optional items
Expanded questions	Particular expanded (only) questions are not covered in the register scope.	Omit these questions

Note: Expansion beyond the basic Core and Expanded questions is suggested only in settings where resources are available and local needs require expansion.

Modification rules There are some fundamental rules that must be observed when making any modifications to the standard Stroke Instrument. These include:

- Never delete a question or measure from the Core (unshaded) Instrument.
 - Never change the standard coding numbers.
 - Place additional questions or measures at the end of the relevant section as an Optional item.
 - Do not place additional questions or measures in between other Core or Expanded (shaded) questions.
 - Code added questions or measures coded with the letter 'X' so they stand out
 - Finally, remove from the Instrument any Expanded sections and Steps (i.e. 2 and or 3) that are not being covered by your site.
 - Send your final draft to the ICS for review **before** starting the register.
-

Continued on next page

Modifying the Stroke Instrument, Continued

Translating the Instrument

Follow the guidelines below to select appropriate translators and ensure accurate and appropriate translation of the Stroke Instrument and all other interviewing materials.

- Initial translation of material should be conducted by at least one translator (ideally by health and survey experts who have a basic understanding of the key concepts).
 - The Instrument must then be back-translated into the original language by another translator to ensure accurate reproduction of meanings (ideally by linguistic experts who can explain the terms used and suggest alternatives).
-

Quality standards for translation

The following are recommended guidelines for translation:

- Translate medical terms into expressions understood by all health professionals.
 - Translate the original intent of the questions with the most appropriate equivalent term in the local language.
 - Develop an inventory of local expressions used as well as comparisons of expressions in other languages.
 - Where there are many dialects and/or languages that are not available in written format, carefully plan specific translation protocols.
-

Registering for Participation

Introduction Once you have addressed the prerequisite actions and identified the scope of your stroke register, you will need to complete a registration form for participation in STEPS Stroke surveillance.

Purpose The purpose of the registration for full participation is to set out:

- Location and health care facilities to be included in the register
- Details about the site coordinator (including expertise)
- Scope of the register and desired goals
- Details of planned overlapping case finding methods which will be used
- Defined population
- Required resources
- Financial support
- Data management environment.
- Contact details.

Registration for participation template A stroke registration form for participation can be found in section 7. Once completed, you will need to forward this to one of the addresses below.

WHO/PAHO/CAREC
STEPS Stroke Surveillance

Dr. Branka Legetic
Pan American Health Organization
525 23rd St., NW
Washington DC 20037

Email: **legeticb@paho.org**

Dr. Glennis Andall-Brereton
Caribbean Epidemiology Centre
16-18 Jamaica Boulevard
Federation Park
Port of Spain
Republic of Trinidad and Tobago

Email: **andallgl@carec.paho.org**

Participation acceptance

Once your registration has been accepted by the ICS, you will be given provisional SSS participation status. Full participation will be granted once you have received ethical approval from your local ethical review committee. You will receive the following from ICS:

Acceptance stage	Received from ICC
Provisional	<ul style="list-style-type: none">• Stroke surveillance site code• Interviewer codes• Hard copy of the Stroke Manual
Full	<ul style="list-style-type: none">• Password to logon to the STEPS stroke web site and download the data entry tool

Getting Ethical Approval

Introduction To ensure that each stroke survey is conducted in a technically and ethically sound manner and in appropriate consideration of the local context, every stroke surveillance registration should undergo ethical review and approval.

Process Ideally, ethical approval should be sought by submission of a proposal and application to a hospital ethics review committee or other relevant body.

Where no such established process exists, it is recommended that an application for ethical review be prepared and submitted through an ad hoc local mechanism within the Ministry of Health.

Making a submission Follow the steps below to make a submission and obtain ethical approval and access to information used as the sampling frame for the survey.

Step	Action
1	Draft a formal submission.
2	Identify and contact the relevant committee, seeking guidance on rules, submission processes and procedures and committee sitting times.
3	Adapt submission as necessary and submit to the appropriate committee requesting guidance on expected timeframe for approval. Note: Emphasize that all data collected are kept confidential.
4	Follow-up with committee to get clearance.

Note: The STEPS stroke coordinating committee can provide further advice on making a submission.

Expected timeframes Preparing and obtaining approval for submissions to ethics committees can take weeks and even months depending on their rules of operation and how often committees sit.

Possible issues Some of the issues that can occur while trying to gain ethical approval include:

- Committee does not sit for months
- Committee takes too long to provide consent
- Ethical approval is declined
- The committee wants modifications to the instrument that threaten its value.

Continued on next page

Getting Ethical Approval, Continued

Informed consent	In addition to getting ethical approval, it is also recommended that there is a process for patients to give verbal and/ or written consent before taking part in the register.
Approaching participants	<p>Important issues to raise in obtaining consent for potential participants or their family members include the following:</p> <ul style="list-style-type: none">• Introducing the institution carrying out the register• Stressing confidentiality• Indicating voluntary nature• Reaching agreement on consent to participate.
Consent letter	A model of a consent letter for patients to give verbal and/ or written consent before taking part in the register is given below. See also section 5, data collection guidelines.



STEPS Stroke Patient Consent

(To be used if 28 day follow-up is included in the stroke register)

Dear patient

Introduction This register is being conducted by (name of country). It is being carried out by professionals from (name of institution).

Confidentiality The information you provide is totally confidential and will not be disclosed to anyone. It will only be used for research purposes. Your name, address, and other personal information will be removed from all records and only a code will be used to connect your response to the register. You may be contacted by the register team again to complete information for the register.

Voluntary participation Your participation is voluntary. If you have any questions about this register you may ask me or contact (name of institution and contact details) or (the site coordinator).

Consent to participate Signing this consent indicates that you understand what will be expected of you and are willing to participate in this register.

Read by Participant		Interviewer	
Agreed		Refused	

Signatures I hereby provide INFORMED CONSENT to take part in the STEPS Stroke surveillance register.

Name: _____ Sign: _____

Next of kin: _____ Sign: _____

Witness: _____ Sign: _____

Section 4: Preparing the Stroke Surveillance Site

Overview

Introduction Once your registration for participation has been accepted and you have received your Stroke surveillance site code (SSS code), you will be able to recruit or obtain staff through secondment and set up the data entry tool.

Intended audience This section is designed for use by those fulfilling the role of the site coordinator.

In this section This section covers the following topics:

Topic	See Page
Recruiting Staff	4-2
Briefing and Training for Data Collection Staff	4-3
Setting up the Stroke Surveillance Site	4-4
Installing and Preparing the Data Entry Tool	4-5
Test Run	4-7

Recruiting Staff

Introduction	<p>The number and type of staff recruited to do data collection will depend on the following:</p> <ul style="list-style-type: none">• Scope and size of the stroke register (including Step 1, 2 and or 3)• Source of data to be collected (ie active recruitment, or retrospective record review)• Qualifications and skills of interested applicants.
Core roles	<p>The core roles of data collection staff are covered in section 2.</p>
Where to get people from	<p>In many countries, recruitment is likely to be an informal process where staff are 'seconded' from other duties within a health facility or health authority. For example, junior staff in training. In this situation, arrangements for their release or scheduled participation may need to be negotiated and explicitly agreed upon.</p> <p>Where there is not sufficient available staff or specific skills are required formal recruitment may be necessary.</p>
Number of staff	<p>For a STEPS Stroke surveillance site that intends to register prospectively 250 cases per year, you will need to recruit two data collection staff.</p> <p>It is normal to screen a much larger number of patients with a range of stroke-like symptoms even vaguely suggestive of a stroke. Data collection must be ensured during sick leave, annual leave, etc.</p> <p>More staff may be needed if data collection is spread over a large area. This includes multiple hospital coverage and in places where death certificates are not centralized.</p>

Briefing and Training for Data Collection Staff

Introduction Training is likely to be informal and will depend on the level of skill and qualifications of data collection staff.

Purpose The purpose of the briefing and training is to ensure:

- Uniform application of the steps stroke surveillance materials
 - Good overall quality of data collected, and
 - Useful and meaningful results reported.
-

What to cover Depending on the qualifications and skills of staff recruited and type of data collection, briefing and/ or training could cover some or all of the following topics:

Topic	What to cover	Reference
STEPS stroke surveillance	<ul style="list-style-type: none">• Basis of STEPS stroke and rationale• About stroke• Key definitions and epidemiological concepts• Key vascular risk factors• Medical treatment	Section 1
Roles	<ul style="list-style-type: none">• Roles and responsibilities of data collection staff	Section 2
Data collection	<ul style="list-style-type: none">• Methods for identifying stroke patients (hot and cold pursuit)• Interview skills• Recording patient responses• Using the STEPS stroke instrument• Using the Q by Q Guide• Administration• Test cases	Section 5 Section 7
Updating the stroke register	<ul style="list-style-type: none">• Using the data entry tool	Section 6

Setting up the Stroke Surveillance Site

Introduction

To set up the STEPS stroke surveillance site, an office space will need to be identified or established for:

- Coordinating the steps stroke register
 - Entering patient data entry into the data entry tool, and
 - Maintaining the stroke register and relevant files.
-

Office equipment and supplies

General office equipment and supplies required for the stroke surveillance site office include:

- Photocopier
 - Shelving
 - Filing cabinets or boxes
 - Telephone
 - At least one computer with internet connection
 - Office stationery supplies
-

Software

The following is a list of software that you will need to have setup on your office computers:

- Microsoft Office '98 or higher recommended for reports, correspondence and general word processing.
- Virus scanning software (if connected to the internet and/ exchanging files outside the office).
- MS Access ('98 or higher) for data entry.
- Standard Data Entry Tool (DET).

For information on installing the DET see page 4-5 and for further information on using the DET see section 6.

Note: The DET is a standard tool that is only available in English.

Other technical requirements

To conduct an "ideal" stroke register, you will need access to brain imaging equipment at the surveillance site. This may not be feasible in all settings.

Installing and Preparing the Data Entry Tool

Introduction

It is important to properly set up and install the Data Entry Tool (DET) prior to starting data collection. The setup process involves:

- creating a folder for the DET
- receiving and installing the DET
- preparing the DET for data entry.

Create DET folders

Follow the steps below to create appropriate STEPS Stroke folders on the computer that will be used to enter data and establish a register.

Step	Action	Recommended folder name
1	In Windows Explorer, create a primary folder (directory) for all your Stroke files, including: <ul style="list-style-type: none">• data• code• documents, and• other files.	Use either: <ul style="list-style-type: none">• C:\SSS200X(insert appropriate year), or• other appropriate drives if your disk is partitioned you are on a network.
2	Create a sub-folder under the STEPS Stroke primary folder to contain your data files	C:\SSS200X\data

Receiving the DET

Once you have been authorized by the ICS to participate in STEPS Stroke, the ICS data analyst will email you the appropriate version of the DET.

Note: Make sure you have ordered the version of the DET which corresponds to the Microsoft Access® version on your computer or local network (e.g. DET for MS Access 97).

Installing the DET

Follow the steps below to install the DET onto your computer.

Step	Action
1	Unzip the DET attachment
2	In Windows Explorer, copy the following files into the C:\SSS200X folder: WHO_Original.mdb WHO_Original_Data.mdb Note: All export files of the Data Entry Tool are automatically stored into this folder.

Continued on next page

Installing and Preparing the Data Entry Tool, Continued

Installing the DET (continued)

Step	Action
3	In Windows Explorer double click the file: WHO_Original.mdb
4	Click OK in the Entering SSS code box.
5	Click on the SSS code button in the configuration window.
6	Enter your SSS code and click OK in the SSS code window.
7	Repeat step 9 to confirm your SSS code.
8	Click the Close button in the configuration window.
9	Click the Close button in the start window.

Error messages You will get one of the following error messages if you incorrectly entered your SSS code and will need to repeat steps 7-10 above to correct:

- invalid data entry
 - wrong sss code
 - invalid SSS code.
-

Test Run

Introduction	<p>Prior to starting the stroke register, it may be useful to complete a test run of the:</p> <ul style="list-style-type: none">• STEPS stroke instrument• case finding process• access to records• case registration process, and• data transfer of results.
Test patients	<p>Identify and approach a sample of 25 patients to be part of the test run. If possible, the test cases should include:</p> <ul style="list-style-type: none">• both men and women• people with differing levels of education• a broad age range (within the target range)• more than one ethnic group (if appropriate).
Feedback	<p>At the end of each interview, ask the patient the following questions and record their feedback:</p> <ul style="list-style-type: none">• Did any of the questions make you feel uncomfortable?• How could we improve the format or layout?• Were there activities that we missed?• How else could we improve this survey?
Evaluation and refining the Instrument	<p>On completion of the trial:</p> <ul style="list-style-type: none">• Compile all patients' comments into a single report.• Where necessary, adapt and refine the Instrument - taking care not to change intended meanings.• Send the instrument to the ICS for comment and quality assurance. <p>Note: The ICS will also provide feedback on the overall quality of collected data.</p>

Section 5: Data Collection Guidelines

Overview

Introduction This section provides generic guidelines for data collection staff.

Intended audience This section is designed for use by those fulfilling the following roles:

- Interviewers
- Stroke principal investigator

In this section This section covers the following topics:

Topic	See Page
Case Finding Methods	5-2
Identifying Stroke Patients in Hospitals (Step 1)	5-3
Identifying Fatal Stroke Patients in the Community (Step 2)	5-5
Estimating Non-Fatal Stroke Events in the Community (Step 3)	5-7
Interview Skills	5-9
Recording Responses for Registration	5-11
Completing the Stroke Instrument	5-12
Guide to Completing the Instrument: All Stroke Events	5-13
Guide to Completing the Instrument: Events Admitted to Hospital (Step 1)	5-15
Guide to Completing the Instrument: Fatal Community Events (Step 2)	5-21
Guide to Completing the Instrument: Non-Fatal Community Events (Step 3)	5-22

Case Finding Methods

Introduction

The main case finding methods used to identify stroke cases are:

- Hot pursuit (active, ongoing recruitment)
 - Cold pursuit (retrospective record review)
 - Combination of hot and cold pursuit.
-

Hot pursuit

Hot pursuit refers to ongoing 'active' identification of all stroke events as they occur. The main purpose is to confirm that the criteria for stroke are met and ensure complete identification of all events including mild stroke events.

Hot pursuit involves regularly checking the following:

- Daily hospital admissions,
 - Hospital separations or discharges
 - Emergency room registers
 - Wards or units
 - Death certificates.
-

Cold pursuit

Cold pursuit refers to retrospective identification of stroke events, for example, based on information from hospital discharge records, or death certificates.

This identification method relies on diagnoses made by several doctors of varying neurological experience who are not working to a protocol. It requires a team identifying and validating stroke events when it is convenient, based on information from routine data sources. Direct examination of the patient is often not possible, and the diagnosis is based on data from records.

Combined approach

Many studies use a combined approach with a mix of hot and cold pursuit to ensure the most complete identification of stroke events (so called overlapping identification sources or overlapping sources of information).

Some of the patients must have been identified as soon as possible after symptoms onset with the possibility of direct examination, while the remaining events are based on routine data.

For example, the researchers have done direct examinations after hospital admission but to ensure the completeness of the data, hospital discharge records, death certificates etc. are checked, physicians are asked to report non-hospitalized stroke events.

Identifying Stroke Patients in Hospitals (Step 1)

Introduction

Surveillance of stroke managed in hospitals should be limited to patients who:

- Are admitted to any unit, ward, division or department of the hospital with a provisional diagnosis of having experienced the onset of a new stroke.
 - In-hospital patients who suffer a stroke due to the treatment of another disease.
-

Identifying stroke patients

Stroke patients may be identified through the following hospital systems and channels:

- Emergency room daybook (or register)
- Admissions book (or register)
- Outpatient clinics
- Radiology departments
- Specialist physicians or neurologists
- Physiotherapists, speech or occupational therapists
- Discharge records
- Death certificates

Note: It is necessary to devise systems in each hospital to detect patients who suffer from an in-hospital stroke, whether intra-operatively or at some other time, and whether in acute or on long-stay wards.

Difficult cases

While many cases are straightforward, stroke has a long differential diagnosis. Resolving the difficult cases requires that the patient be assessed by an experienced medical practitioner and preferably by an internal physician or a specialist neurologist.

Re-assessment of the patient at least 24 hours after the initial presentation may be vital to differentiate stroke from TIA and other neurological or medical diseases such as hemiplegic migraine and epilepsy.

Death following stroke

In order to assess trends in case fatality, a system for accessing details on all deaths occurring in stroke patients registered is necessary. Date of death should be registered in the Instrument, together with details of cause of death. Vital status of all patients should be known at 28 days after the onset of stroke. This length of follow up is not always feasible. A minimum follow up is 7 days.

Continued on next page

Identifying Stroke Patients in Hospitals (Step 1) Continued

Diagnostic criteria

Each registered stroke event must meet the standard WHO clinical definition of stroke (see p1-4). Key features of the clinical definition are as follows:

- Sudden onset
- Neurological deficit
- Lasting 24 hours or longer
- Of presumed vascular origin.

The table below provides an example of some of the diagnoses that should be considered for STEPS stroke registration.

Stroke specific	Focal and global signs that could be caused by stroke
<ul style="list-style-type: none">• (Acute) stroke <i>or</i> (acute) cerebrovascular episode• Cerebral <i>or</i> cerebellar embolus, thrombosis <i>or</i> infarction• Occlusion, thrombosis <i>or</i> embolus of carotid, (pre) cerebral <i>or</i> vertebral artery• Lacunar hemorrhage <i>or</i> stroke• Subarachnoid, (primary) intracerebral, cerebellar <i>or</i> pontine hemorrhage <i>or</i> stroke• Transient (cerebral) ischemic attack	<ul style="list-style-type: none">• (Acute) hemiplegia <i>or</i> (acute) hemiparesis• Faint, fit, funny turn, (acute) confusional state• Loss of consciousness• (Acute) dysphasia, dysarthria, dyspraxia• Homonymous hemianopia• Amaurosis fugax• Acute monocular blindness

Note: Further details on symptoms for the three major stroke types can be found in section 1, About Stroke.

Residency criteria

If a population based register, the stroke case, to be eligible, must be resident in the defined population at the time of the onset of the stroke.

Identifying Fatal Stroke Patients in the Community (Step 2)

Introduction

The three main methods for identifying and estimating the number of stroke patients that die from a stroke, but who do not reach hospital facilities:

- Death certificates
 - Verbal autopsies
 - Medical autopsy.
-

Death certificates

Communities that have routine medical certification of cause of death can provide direct data on deaths due to stroke. Note that delays in processing death registrations and certificates may occur and also a wide variety of terms may be used to describe fatal stroke. When in doubt, verification or follow up is essential. Methods for searching death registrations may include:

- Electronic keyword search
 - Manual record search by visual sighting.
-

Verbal autopsies

Verbal autopsies (VAs) are increasingly being used to monitor the distribution of deaths by cause in places where medical certification of cause of death is uncommon. This technique is based on the assumption that most causes of death have distinct symptom complexes, and that these can be recognized, remembered and reported by health professionals or lay respondents.

Official WHO verbal autopsy for adult deaths is currently being developed. A provisional form for assisting with the process of verifying deaths which may possibly have been due to stroke is available on request from the ICS for those SSS planning population based registers.

Follow up with this modified VA is essential where cause of death is cited in such vague terms as:

- Ill defined
- Unknown
- "old age"
- Senility.

Sensitivity in obtaining this information must be adhered to at all times.

Continued on next page

Identifying Fatal Stroke Patients in the Community (Step 2), Continued

Verbal verification

When interviewing health professionals or family members about the signs and symptoms associated with a possible stroke event, the following questions provide a structure to the interview:

- Was the deceased ill prior to death?
- Did s/he have weakness on one side of the body prior to death?
- Did that weakness develop suddenly?
- Did it last more than 24 hours?
- Was it a sudden death (died within 24 hours)?
- Was there a history of severe headache just prior to death?
- How many days was the patient ill before death?
- Was the patient seen by a medical or health professional?
- Was the patient admitted to hospital or a clinic? How many nights?

A detailed description of the previous illnesses, treatment, and events leading to death should be prepared and used in the decision by the site coordinator as to whether the criteria for stroke have been met.

Validation of codes and diagnosis

Both the codes used and diagnosis of stroke as the immediate or underlying cause of death should be validated as indicated in the table below.

Validation of	Based on
Codes	<ul style="list-style-type: none">• Medical and medico-legal records (within 28 days of death).• Interview with decedent's next-of-kin or other informant (verbal autopsy).
Diagnosis	<ul style="list-style-type: none">• Clinical signs according to the stroke definition• Neuro imaging or autopsies.

Medical autopsies

Since medical autopsy rates are declining in many countries autopsies are unlikely to provide a substantial coverage of fatal strokes. However, records of post mortem examinations are an accessible way of getting information for the surveillance system. They provide a valid diagnosis, and contribute to a more complete understanding of the stroke occurrence in the defined population.

Estimating Non-Fatal Stroke Events in the Community (Step 3)

Introduction

The main methods for estimating numbers of non-fatal events in the community include:

- Tracking local medical practices and health facilities by survey
 - Hemiplegia/ hemiparesis survey (prevalence survey)
-

Primary health care facilities

Where general practitioners are widely used at primary health care facilities, these should be included as part of the case finding methods.

In some countries there are only a few general practitioners or only a proportion of stroke patients who ever have contact with them. In these sites, local healers may be the primary contact person and it is important to consider their potential for collaboration.

General practitioners

You will need to use different survey techniques depending on the size of the defined population to determine the number of general practitioners to include, as indicated in the table below:

If the population is	Then...
Small (limited size)	Include all the general practitioners and local health facilities in the register (eg. public health care centres, nursing homes, rehabilitation centres etc).
Large (entire population)	Survey a representative sample of medical practitioners to assess the number of cases that they have managed over a defined, preceding period.

Local healers

Given instructions on stroke symptoms, local informal healers may be able to provide a contact to the patient, who then can be examined for stroke symptoms.

Note: This procedure is likely to underestimate the true rate as mild cases are unlikely to be detected, but the overall effect on the estimates is likely to be minor.

Continued on next page

Estimating Non-Fatal Stroke Events in the Community (Step 3), Continued

Sampling community based facilities

Informing the community about the purposes of a stroke surveillance system is crucial in enlisting assistance in case finding for the hard to reach people who are not listed with a registered facility or hospital.

Identification of these patients involves the collaboration and cooperation of general practitioners and other health care providers in the community, but their identification is vital for the accurate determination of stroke incidence.

These strokes are a combination of milder and more severe strokes than those that present to hospital, and consequently their inclusion influences case fatality ratios. One approach is to notify all general practitioners (or equivalent) of the register. They should be provided with information kits and memory aids to focus their attention on the register and to alert the team to any cases in their practice who have a stroke- especially if not admitted to a hospital.

In addition to general practitioners (or equivalent), the team will need to maintain close liaison with community health nurses and village elders/church leaders to ensure ongoing support and referral of potentially eligible cases to the register.

Prevalence of stroke

Hemiplegia/ hemiparesis survey

One way of estimating the proportion of non fatal stroke events is to obtain an estimate from the prevalence of survivors. In most communities the causes of adult-onset hemiplegia or hemiparesis are limited to stroke and head injury and can be distinguished from patient history.

If the incidence of residual hemiplegia following stroke and the survival time are constant within a given community, trends in the prevalence of hemiplegia will reflect trends in the incidence of stroke.

This could be useful for stroke surveillance because hemiplegia is recognisable and identifying cases does not require self diagnosis. The prevalence of hemiplegia can therefore be identified by questionnaire based population surveys or interviews with a representative from selected households. The problem, however, is that even prevalence of stroke is relatively rare.

If such a survey has already been undertaken, check to see if information was obtained on the proportion who said they had never been admitted to a hospital facility. This could act as a proxy measure of the non fatal community based group.

Notes: The linkage between prevalence of hemiplegia/ hemiparesis and incidence of stroke has not been validated in a study so far.

Interview Skills

Introduction Although much of the data that needs to be collected can be obtained from records, some contact with patients or next of kin may be required.

Participation The patient (or person being interviewed) needs to feel comfortable about the interview and can refuse to be interviewed as participation is voluntary. The interview should therefore be as natural as possible and conducted politely, like a normal conversation.

Behaviour and tact The table below provides guidelines on appropriate behaviour during an interview:

Behaviour	Guidelines
Respect confidentiality	Maintain the confidentiality of all information you collect.
Respect patient's time	You are asking patients for their time so be polite and prepared to explain.
Tact	If you feel that a person is not ready to assist you, do not force them but offer to come back later.
Friendly disposition	Act as though you expect to receive friendly co-operation and behave accordingly.
Body language	Maintain good eye contact and adopt appropriate body language.
Pace of interview	Don't rush the interview. Allow the patient enough time to understand and answer a question. If pressured, a patient may answer with anything that crosses their mind.
Patience	Be patient and polite at all times during the interview and ensure you have set aside enough time for patients with aphasic disturbances.
Appreciation	Thank them for their help and cooperation.

Continued on next page

Interview Skills, Continued

Handling refusals

Be prepared to obtain cooperation from a patient who does not want to be interviewed. In general, be pleasant, good-natured, and professional and most patients will cooperate.

If...	Then...
The patient becomes defensive	<ul style="list-style-type: none">• Show patience and understanding• Provide token agreement and understanding of his/her viewpoint, that is, saying something like, 'I can understand that' or 'You certainly have the right to feel that way.'• Convey the importance of the register to the patient and that all stroke patients are being registered.
You may have visited at a bad time.	Try again later.
The patient may have misunderstood the purpose of the visit.	Try to explain the purpose again.
You think you may get a 'no'	Try to leave and suggest coming back later before you get a partial or an absolute 'no'.

Patient consent

Each patient (or family member) should provide verbal and/or written consent in accordance with local standards before taking part in the register.

A sample patient consent form is discussed in section 3-15.

Recording Responses for Registration

Introduction

All results that are recorded on the STEPS instrument must be written as clearly as possible to avoid ambiguity and confusion when checking and entering the results.

Requirements

Some general requirements for recording survey information are as follows:

- Record the patient identification number on every page of each instrument.
 - Do not erase any notes made.
 - If a question has been skipped by mistake, correct it.
 - If a patient changes his/her mind on one of the options, record the new answer.
 - Record only answers that are relevant to the register.
 - Record comments or explanations in brackets in the Instrument next to the corresponding question.
 - Do not get too absorbed recording. Keep the patient's interest by saying the patient's response aloud as you write it down.
 - Standard agreement on how to write numbers.
-

Handling issues

Use the table below to help with some common issues you may encounter.

If...	Then...
You are uncertain about a response	Repeat the question and record the answer exactly. Do not paraphrase a response.
A question doesn't apply or the patient doesn't know and these options are not available on the Instrument	For "don't know" record: 9, 99 or 999 etc.
You have missed a question	Go back and ask the question, making a note that the question was asked out of sequence.

Checking and editing

At the end of each interview check the Instrument and make sure that:

- All the questions have been answered.
 - The information recorded is clear and legible for others to read.
 - Probing comments are indicated.
 - Check that all the information has been completed including the ID number on every page.
 - Review the Instrument to check it is complete and that every question has been answered.
-

Completing the Stroke Instrument

Introduction	<p>Once the standard stroke Instrument has been translated and printed it is ready for use during the register.</p> <p>One Instrument is to be completed for every eligible stroke event. Note: one person may have more than one stroke event over the year of observation. All items on the Instrument must be completed for the response to be valid.</p>
Cover page	The bottom part of the first page of the Instrument contains identification information, including the patient names. It is very important that these details are kept confidential at all times and that you tell the patient that they will be kept confidential.
Core and core expanded items	The Instrument contains Core (un-shaded) and Expanded (shaded) response options for each Step you will need to complete.
Introductory statements	Where a section of items has an introductory statement, you must read this out to the patient.
Entering the patient's response	For some items on the Instrument, there may be one or more possible responses. Each possible response has an associated code. You will need to enter the appropriate response code in the box for each item. For example:

Stroke classification

(S1 6)	What subtype of stroke was diagnosed? <i>[select one]</i>	Ischemic stroke	(1)	[2]
		Intracerebral hemorrhage	(2)	
		Subarachnoid hemorrhage	(3)	
		Unspecified type	(4)	

Unknown responses The table below shows what to enter as a last resort where the patient does not respond with a standard response.

If a hospital record or patient response is	And number of [] is	Then enter
Unknown Don't know	[]	9
Unknown Don't know	[] []	9 9
Unknown Don't know	[] [] []	9 9 9

Continued on next page

Guide to Completing the Instrument: All Stroke Events

Introduction	Guidelines on how to complete some questions in the All Stroke Events section of the Instrument are given below, with further guidance given in the Question by Question Guide in section 7.
Identification number	The patient identification number is to be written in the boxes at the top of each page of the Instrument and all patient specific documents at the time the completed Instrument is being entered into the register.
Patient identification and patient characteristics I 1 - I 3 (Core)	<p>Accurate core participation and patient characteristic information is essential for analysing and reporting on the overall results of the STEPS Stroke surveillance.</p> <p>Patient identification and patient characteristics should be completed for every patient documented in Step 1, Step 2 and Step 3. If the age and sex of a patient has been missed out, the Instrument cannot be used in the analysis, as most analyses are grouped by these criteria.</p>
Contact name and address I 9 - I 13	An acute stroke event often results in dramatic consequences for the patient after discharge from hospital. This may mean the patient goes to live with relatives or a nursing home for long term care. The contact person, phone number and address should therefore be for someone who knows about the actual living situation of the patient. Children or other close relatives could serve as contact persons for the patient. The relationship of the contact person to the patient should also be documented.
Dates of birth and age I 14 (Core)	In some countries exact dates of birth and/or age are not known. In these situations age has to be estimated. To estimate someone's age, you will need to ask them how old, or at what stage in life they were at the time that a number of widely known major local events occurred.
Information on acute stroke event I 20 - I 22 (Core)	<p>If the exact onset of stroke symptoms is unknown (e.g. stroke occurred during sleeping), ask the patient or another person when the first symptoms of stroke were noticed and enter that date.</p> <p>To differentiate between a first-time event and a recurrent event it is important to obtain information about possible previous strokes. Please note that the following are not counted as a stroke:</p> <ul style="list-style-type: none">• previous TIA• silent strokes (ie detected by scanning but did not result in a neurological deficit longer than 24 hours).

Continued on next page

Guide to Completing the Instrument: All Stroke Events

Continued

Expanded items Expanded questions are shown in the shaded boxes. Some of these questions may have been adapted so the terms and phrases make sense to patients in your environment. Some of the adaptations may include relevant:

- Ethnic, racial and or cultural groups
- Highest level of education
- Categories of work
- Income level.

Choosing expanded items Each site coordinator should choose which of the expanded items will be included in the final Instrument. It is a question of balance between obtaining the minimum data (core items) and what can reasonably added without additional effort and cost.

Choosing expanded questions depends on local circumstances and the use to which the information will be put. As with core questions, expanded questions should not be altered.

Guide to Completing the Instrument: Events Admitted to Hospital (Step 1)

Introduction

This section is to be completed for all eligible stroke patients admitted to hospital. Information collected includes:

- Hospital admission
- Stroke classification
- Vascular risk factors
- Medical treatment
- Secondary prevention
- In-hospital management
- Follow up of the patients

Note: Each of these is explained in more detail below.

Hospital admission S1 1 (Core)

Stroke patients admitted to hospital must have survived until hospitalization, and must have been able to get to the hospital either:

- by themselves
- with the help from relatives/care givers, or
- using any kind of emergency medical service.

Note: Despite differences between countries and changes in admission practices over time, data based on hospitalized events gives valuable information for local health authorities, and constitutes the first step to a better understanding of stroke in the population.

Hospital departments S1 2(Expanded)

There are seven possible answers for indicating in which departments or units the patient was treated. The available options are explained in the table below.

Department/unit	Refers to patients managed at..
Intensive care	An intensive care unit, including any type of acute medical unit.
Medical	A general medical ward, including a geriatric unit.
Neurological	A general neurological ward.
Neurosurgical	A general neurosurgical ward.
Rehabilitation	A specialized rehabilitation unit, except a rehabilitation stroke unit.
Stroke	Acute and rehabilitation stroke units.
Other	Other units, e.g. outliers or patients on surgical wards.

Continued on next page

Guide to Completing the Instrument: Events Admitted to Hospital (Step 1), Continued

Living situation
S1 3 (Core) Living condition options are explained in the table below.

Option	Refers to patients living
Independent at home	Without depending on any assistance from relatives or professionals
Dependent at home	Depending on assistance from relatives or professionals
Community facility	In nursing or residential homes, serviced flat or other long term care facility.

Modified Rankin scale
S1 4 (Expanded)

If possible, the Modified Rankin scale prior to acute stroke event should be assessed retrospectively based on the information provided by patient and/ or close relatives. The number corresponding to the patient's functional level is to be entered. The scale is divided into 6 levels (from level 0 to level 5) as described in the table below.

Scale		Description
0	No symptoms	No symptoms at all
1	No significant disability	No significant disability despite symptoms, ie. can do all usual activities
2	Slight disability	Unable to do all previous activities, but able to look after own affairs without assistance
3	Moderate disability Able to walk without assistance	Requiring some help but able to walk without assistance
4	Moderate disability Unable to walk without assistance	Unable to walk without assistance, and unable to attend to won bodily needs without assistance
5	Severe disability	Bedridden, incontinent, and requiring constant nursing care and attention.

Note: The modified Rankin Scale measures independence rather than performance of specific tasks. Mental as well as physical adaptations to the neurological deficits are incorporated, and the score gives an impression of whether the patients can look after themselves in daily life.

Continued on next page

Guide to Completing the Instrument: Events Admitted to Hospital (Step 1), Continued

Neurological signs S 1 5 (Core)

Neurological deficits, for example, disturbances of consciousness, are an important predictor of stroke severity. Neurological deficits present at the first medical examination after hospitalization should be documented to adjust potential differences in outcome and disability for stroke severity. The different levels of deficit are explained in the table below.

Neurological deficit type	Refers to
Disturbed consciousness	Disturbances of consciousness, including semi consciousness, e.g. not fully aroused, and coma, either response to pain only or no response at all
Weakness/paresis	Motor deficits of the upper or lower limbs.
Speech disturbances	Speech disturbances present on admission, like aphasia or dysarthria.

Stroke classification S1 6 (Core)

Stroke events can be classified into either

- ischemic stroke
- intracerebral haemorrhage
- subarachnoid haemorrhage, or
- unspecified.

It is recommended that stroke types are classified as a result of neuro-imaging.

Whether an event is haemorrhagic versus ischemic is also of importance from a clinical perspective in terms of treatment and early secondary prevention, as aspirin should not be given to patients with haemorrhagic stroke and anticoagulation as well as thrombolysis is obviously contraindicated in hemorrhagic strokes.

Where no diagnostic examination was done to verify the subtype of stroke, choose the option *Unspecified*.

Note: For further details on stroke classification, see section 1-4, About Stroke.

Continued on next page

Guide to Completing the Instrument: Events Admitted to Hospital (Step 1), Continued

Subtype diagnosis S1 7 (Core)

Diagnosis of stroke subtype, refers to patients where the subtype classification was verified from one of two methods as follows:

Diagnosis by	Explanation
Clinical diagnosis alone	Clinical diagnosis alone and was not verified by brain imaging (or in subarachnoid hemorrhage on lumbar puncture) in non-fatal cases or also by medical autopsy in fatal cases; please indicate also clinical diagnosis alone if any scoring system not based on brain imaging or medical autopsy was used
Diagnostic techniques	In non-fatal cases to patients where the subtype of stroke was verified by brain imaging; subtype verification of subarachnoid hemorrhage might also be based on lumbar puncture alone; in fatal cases verification of stroke subtype might also be based on medical autopsy.

Risk factors S1 10 (Expanded)

The main modifiable risk factors that are present pre-stroke are listed and defined in the table below.

Risk factor	Defined as a patient who pre-stroke..
Atrial fibrillation	Has atrial fibrillation in ECG prior to stroke (records seen) or during hospitalization.
Current tobacco use	<ul style="list-style-type: none"> Is a current tobacco user (smoking and other forms of tobacco), or Was a recent tobacco user but stopped less than 3 months before acute stroke event.
Diabetes mellitus	<ul style="list-style-type: none"> Has been diagnosed with or has self reported diabetes mellitus, and Uses antidiabetic drugs
Hypercholesterolemia	<ul style="list-style-type: none"> Has reported elevated plasma total or LDL cholesterol level, or Uses lipid-lowering medication
Raised blood pressure	<ul style="list-style-type: none"> Has diagnosed or self reported raised blood pressure, or Uses antihypertensive drugs.

Continued on next page

Guide to Completing the Instrument: Events Admitted to Hospital (Step 1), Continued

Pharmaceutical treatment S1 11 S1 12 (Core)

Pharmaceutical treatment means continuous medication. The only exception is for thrombolysis, which is only given one time. The table below lists the categories of drug type and drugs used in each category.

Drug type	Including
Anticoagulant	<ul style="list-style-type: none"> • Warfarin • Heparin
Anti diabetic	<ul style="list-style-type: none"> • Antidiabetic medications • Insulin injections
Antiplatelet	<ul style="list-style-type: none"> • Aspirin • Clopidogrel • Dipyridamol
Cholesterol lowering	<ul style="list-style-type: none"> • Statins
Blood pressure lowering	<ul style="list-style-type: none"> • Thiazides • Angio-tensin targeting agents • Beta-blockers • Calcium channel blockers

In hospital assessment S1 13 - 14 (Expanded)

The in-hospital assessment questions refer to assessments of the listed disorders during hospitalization, irrespective of whether the patient was treated or not after the first visit.

Patient discharge S1 15 - 18 Core)

If the patient is alive at discharge (S1 18), there are three possible destinations. These are explained in the table below.

Option		Refers to patients discharged to
1	Home	Private address (either the same or a new address)
2	Other hospital	<ul style="list-style-type: none"> • Another hospital • Rehabilitation unit • Rehabilitation hospital • Long-stay hospital
3	Community facility	Facilities with access to service and staff eg: <ul style="list-style-type: none"> • Nursing or residential homes • Long term care facilities for psychiatric disorders • Serviced flat, or • Assisted living

Continued on next page

Guide to Completing the Instrument: Events Admitted to Hospital (Step 1) Continued

Modified Rankin scale S1 9 (Expanded)

If the patient is alive at discharge, the Modified Rankin scale should be assessed just before discharge from hospital. The number corresponding to the patient's functional level is to be entered. The scale is described on page 5-16.

Follow up at day 28 F1 - F7 (Optional)

Follow up on day 28 (from onset of stroke) provides valuable information about the long-term burden of stroke. These optional questions may be difficult to obtain for all registered patients. If a patient or contact person could not be contacted on day 28, try to get all necessary information as soon as possible, and within the next few days.

Some possible ways to follow up with patients after discharge from hospital include:

- Direct examination, e.g. during a home visit, outpatient department or in hospital.
- Medical record review, if the patient is still in the hospital at day 28.
- Telephone interview with the patient or close relative.
- Questionnaire posted to the patient.

Note: Confidentiality, ethical issues and other legal aspects in terms of performing a follow up should be clarified before starting data collection.

Guide to Completing the Instrument: Fatal Community Events (Step 2)

Introduction

Step 2 covers identifying and registering every fatal stroke event treated in community and not admitted to hospital.

How information is collected S2 3 (Core)

There are three main methods for collecting information about fatal stroke events in the community. These include:

- verbal autopsy
- death certificates
- medical autopsy.

For further information on each of these methods see page 5-5.

International classification of diseases (ICD) S2 4 - 5 (Core)

The International Classification of Diseases (ICD) system is commonly used to record the cause of death on death certificates. There are three versions of the ICD codes and a range of eight or nine coded diseases that may relate to stroke as the cause of death. Some of these diseases will not meet the definition of stroke, but should be included in all broad searches for stroke events.

The ICD versions and codes are as follows:

Version	Codes	Disease
ICD 9	431	Intracerebral haemorrhage
	432	Other and unspecified intracranial haemorrhage
	433	Occlusion and stenosis of precerebral arteries
	434	Occlusion of cerebral arteries
	435	Transient cerebral ischemia
	436	Acute but ill-defined cerebrovascular disease
	437	Other Ill-defined cerebrovascular disease
	438	Late effects of cerebrovascular disease
ICD 10	I60	Subarachnoid haemorrhage
	I61	Intracerebral haemorrhage
	I62	Other non-traumatic intracranial haemorrhage
	I63	Cerebral infarction
	I64	Stroke, not specified as haemorrhage or infarction
	I65	Occlusion and stenosis of precerebral arteries, not resulting from cerebral infarction
	I65	Occlusion and stenosis of cerebral arteries, not resulting from cerebral infarction
	I67	Other cerebrovascular diseases
	I68	Cerebrovascular disorders in diseases classified elsewhere
	I69	Sequelae of cerebrovascular disease

Continued on next page

Guide to Completing Step 2: Fatal Community Events, Continued

Verbal autopsies (Optional)

The purpose of Verbal autopsies (VA) is to describe cause of mortality at a community or population level where no better alternative resources exist. Verbal Autopsies are based on interviews with friends and relatives of a deceased person. After an interview has been conducted, the following takes place:

- A panel of physicians reviews the forms and assigns a probable cause of death.
- Medical records coders trained in ICD rules select and code the underlying cause of death, according to a code score.
- Mortality results are tabulated using a standard list capable of generating comparable mortality statistics.

Unfortunately, the tools and methods employed are often imperfect and require rigorous validation and continuous quality assurance.

Guide to Completing the Instrument: Non-Fatal Community Events (Step 3)

Introduction

Step 3 covers identifying and estimating the non-fatal stroke events treated in community and not admitted to hospital. This is the most difficult component of case finding, and efforts at estimating the number of people who would otherwise be missed by a focus only on patients admitted to hospital are essential to show the true incidence of stroke.

How information is collected S3

There are two main methods for estimating non-fatal stroke events in the community. These include:

- Tracking medical practices (health facilities) by survey, and
- Prevalence or Hemiplegia/ hemiparesis survey.

For further information on each of these methods see page 5-7.

Other methods such as capture/re-capture using multiple overlapping sources are available. Please discuss approaches relevant in your setting with the ICS.

Section 6: Data Entry and Data Management

Overview

Introduction This section covers all the tasks that need to be conducted to enter and manage the STEPS Stroke data in the Data Entry Tool (DET) to gradually build up a register.

Intended audience This section is designed for use by those fulfilling the following roles:

- Data collection staff
- Principal investigator

In this section This section covers the following topics:

Topic	See Page
Data Entry	6-2
Data Management	6-5
Creating Reports	6-7
Exporting Data	6-8

Data Entry

Introduction

STEPS Stroke surveillance data from completed STEPS stroke instruments is to be entered into the data entry tool by trained data collection staff.

Data entry process

Data entry is a systematic process that covers the following main stages:

Stage	Description
1	Entering new patient data.
2	Entering the Identification number on patient instruments.
3	Validation and error correction.
3	Backing up.
4	Storing and filing the instruments.

Opening the DET

Follow the steps below to open the DET start window.

Step	Action														
1	Open the DET program by clicking the WHO_Original.mdb file in Windows Explorer.														
2	The start window will appear. The function of each button is explained in the table below. <table><tr><th>Click the button</th><th>To</th></tr><tr><td>New Patient</td><td>Enter new patient records</td></tr><tr><td>Search</td><td>Search for entered data</td></tr><tr><td>Reports</td><td>Generate reports of entered data</td></tr><tr><td>Data Export</td><td>Export entered data</td></tr><tr><td>Delete Patient</td><td>Delete entered data</td></tr><tr><td>Close</td><td>Close the data entry tool (DET)</td></tr></table>	Click the button	To	New Patient	Enter new patient records	Search	Search for entered data	Reports	Generate reports of entered data	Data Export	Export entered data	Delete Patient	Delete entered data	Close	Close the data entry tool (DET)
Click the button	To														
New Patient	Enter new patient records														
Search	Search for entered data														
Reports	Generate reports of entered data														
Data Export	Export entered data														
Delete Patient	Delete entered data														
Close	Close the data entry tool (DET)														

Enter all stroke events data

Follow the steps below to enter new patient data from the All Stroke Events section of the completed STEPS stroke instrument.

Step	Action
1	Click the New Patient button in the start window.
2	A unique identification number for each patient will be generated by the Data Entry Tool. Note: This consists of the joint SSS code (5 digits) and the patient ID code (6 digits).

Continued on next page

Data Entry, Continued

Enter all stroke events data (continued)

Step	Action				
3	<p>Write the whole number in the Identification Number boxes at the top of each page of the patient's completed STEPS Stroke Instrument as follows:</p> <p style="text-align: center;"> SSS Code Patient ID </p> <p style="text-align: center;"> Identification Number [1][1][2][2][0] [0][0][0][0][0][1] </p>				
4	<p>Write the first 5 digits (the SSS code) of this number on page 1, I 1 of the paper copy of each patient's completed STEPS Stroke Instrument as follows:</p> <p>Patient Identification and Patient Characteristics (I 1) Stroke Surveillance Code [1] [1] [2] [2] [0]</p>				
5	When you have recorded the number, Click Ok				
6	Enter data into the All Stroke Events window exactly as it is written on the paper Instrument.				
7	<table border="1"> <thead> <tr> <th>If</th><th>Then</th></tr> </thead> <tbody> <tr> <td>Expanded items (grey boxes on the instrument) have been completed on the instrument</td><td>Click the long grey buttons at the bottom of each window to enter expanded data.</td></tr> </tbody> </table>	If	Then	Expanded items (grey boxes on the instrument) have been completed on the instrument	Click the long grey buttons at the bottom of each window to enter expanded data.
If	Then				
Expanded items (grey boxes on the instrument) have been completed on the instrument	Click the long grey buttons at the bottom of each window to enter expanded data.				
8	Click the Next button to move to the next window, or Back to edit a previous window.				
9	<p>Log in a spreadsheet or notebook all discrepancies, questions and problems (irregularities) that you cannot resolve. Include:</p> <ul style="list-style-type: none"> • patient id number • instrument code number (eg. i 14) • comment. 				
10	Click close when you have completed entering the All Stroke Events data.				

Continued on next page

Data Entry, Continued

Entering Step 1, Step 2 and Step 3 data

After entering the All Stroke Events data, you will come to the Selection window.

Step	Description								
1	Click the appropriate button to enter each Step as follows. <table><tr><th>Step</th><th>Description</th></tr><tr><td>1</td><td>Events admitted to hospital</td></tr><tr><td>2</td><td>Fatal events in the community</td></tr><tr><td>3</td><td>Non-fatal events in the community</td></tr></table>	Step	Description	1	Events admitted to hospital	2	Fatal events in the community	3	Non-fatal events in the community
Step	Description								
1	Events admitted to hospital								
2	Fatal events in the community								
3	Non-fatal events in the community								
2	When data entry is complete, click the Finish button								

Validation and error correction

Before moving to the next patient Instrument, check and resolve any inconsistencies and/or errors noted in the log book or spreadsheet.

Backing up data

The computer used for data entry should be backed up at the end of each week.

Filing the instruments

All completed Instruments that have been entered into the DET should be marked 'entered' on the front page and filed in a secure location.

Data Management

Introduction

To manage the STEPS Stroke data entered using the DET, you may need to perform the following functions:

- search for a patient record
- edit data, and
- delete a patient record.

Search for a patient record

Follow the steps below to search for a patient record.

Step	Action
1	Launch the Data Entry Tool from Windows Explorer
2	Click the Search button from the Start window
3	To find a patient record, either enter the patient: <ul style="list-style-type: none">• ID number (last 6 digits or the Identification number), or• Family name and /or• First name
A successful search by ID opens the Selection window.	
A successful search by patient's name opens the Register window where all matches for the entered name are listed.	
4	If you searched by patient name, highlight the ID of the correct patient name and click Go to Patient.

Find and edit data

Follow the steps below to find and edit specific patient data.

Step	Action
1	Select the appropriate button corresponding to the section of the instrument you want to search in the selection window, e.g. <ul style="list-style-type: none">• All stroke events• Step 1• Step 2• Step 3
2	Choose the Next and Back buttons to find the specific data.
3	Edit the data and close.

Continued on next page

Data Management, Continued

Deleting a patient record

Follow the steps below to delete a patient record.

Step	Action
1	Launch the Data Entry Tool from Windows Explorer
2	Click the Delete Patient button from the Start window
3	Enter a patient ID and click the Search button.
4	Click Yes to delete the patient record.

Creating Reports

Introduction

You can create and print the following reports from patient data entered into the register using the data entry tool:

- sex and age distribution
 - stroke subtype distribution.
-

To create a report

Follow the steps below to create a report.

Step	Action
1	Ensure your printer is connected and on.
2	Launch the Data Entry Tool from Windows Explorer.
3	Click the Reports button from the Start window.
4	Click the type of report you from the Reports window.
5	The selected report will be automatically printed to your printer.

Note: Age distribution (stratified by stroke subtype) can only be calculated after you have created an export file. See Exporting data on page 6-8.

Exporting Data

Introduction

To calculate age distribution (stratified by stroke subtype) or to export the data to other software for statistical analysis, you will have to create an export file.

Procedure

Follow the steps below to create an export file:

Step	Action						
1	Launch the Data Entry Tool from Windows Explorer.						
2	Click the Data Export button from the Start window.						
3	<table><tr><th>Click the button</th><th>To automatically</th></tr><tr><td>Complete</td><td>Create the following text and Excel files of the complete data:<ul style="list-style-type: none">• AcompleteTab.txt• CompleteTab.xls</td></tr><tr><td>Anonymous</td><td>Make the data anonymous by removing identification details data and create the following text or Excel files for data transfer:<ul style="list-style-type: none">• AnonymTab.txt• AnonymTab.xls</td></tr></table>	Click the button	To automatically	Complete	Create the following text and Excel files of the complete data: <ul style="list-style-type: none">• AcompleteTab.txt• CompleteTab.xls	Anonymous	Make the data anonymous by removing identification details data and create the following text or Excel files for data transfer: <ul style="list-style-type: none">• AnonymTab.txt• AnonymTab.xls
Click the button	To automatically						
Complete	Create the following text and Excel files of the complete data: <ul style="list-style-type: none">• AcompleteTab.txt• CompleteTab.xls						
Anonymous	Make the data anonymous by removing identification details data and create the following text or Excel files for data transfer: <ul style="list-style-type: none">• AnonymTab.txt• AnonymTab.xls						
4	Click Close to return to the Start window.						

WHO STEPS STROKE INSTRUMENT

<INSERT COUNTRY/SITE NAME>

All Stroke Events

For further guidance on All Stroke Events, see Section 5, page 5-15

Patient Identification and Patient Characteristics

- | | | |
|-------|--|---|
| (I 1) | Stroke Surveillance Site Code | [] [] [] [] [] [] |
| | <i>Insert 1st 5 digits of automatically generated code from DET</i> | |
| (I 2) | Interviewer Code | [] [] [] [] |
| | <i>Insert code provided by the ICS</i> | |
| (I 3) | Date of completion of the instrument | [] [] [] / [] [] [] / [] [] [] [] [] [] |
| | | d d m m y y y y |

Patient individual records

- | | | |
|-------|---|-------------------|
| (I 4) | Patient's family name
<i>Use CAPITALS, include all names</i> | [_____] |
| (I 5) | Patient's first name
<i>Use CAPITALS, include all names</i> | [_____] |
| (I 6) | Contact phone number
<i>Include area codes (optional)</i> | [_____] |
| (I 7) | Contact address
<i>For follow-up questionnaires (optional)</i> | [_____
_____] |
| (I 8) | Resident? | [] []
YES NO |
| (I 9) | Unique identification number where available
<i>number, PID etc (optional)</i> | [_____] |

Contact person of patient

Include contact person who can confirm the living situation of the patient

- | | | |
|--------|---|------------------|
| (I 10) | Contact person's family name | [_____] |
| (I 11) | Contact person's first name | [_____] |
| (I 12) | Contact person's phone number | [_____] |
| (I 13) | Contact person's address | [_____
_____] |
| (I 14) | Relationship of contact person to the patient | [_____] |

Demographic characteristics

- (I 15) Date of birth [] [] / [] [] / [] [] [] []
If date of birth is unknown, enter age [] [] [] d d m m y y y y
- (I 16) Sex Male (1) []

[select one]

Female

(2)

- | | | | |
|--------|---|-------|---------|
| (I 17) | What is your <i>[insert relevant ethnic/racial groups defined according to local demographic needs]</i> | XX | (1) [] |
| | | XX | (2) |
| | | XX | (3) |
| (I 18) | If other ethnicity, please state [_____] | Other | (4) |

Socioeconomic status

- (I 19) What is the highest level of education the person has completed?
[select one]
*If a person attended a few months of the first year of secondary school but did not complete the year, record "primary school completed".
If a person only attended a few years of primary school or never went to school, record "no formal schooling"*
(Insert country specific education level list)

(I 20) Which of the following best describes the main work status of the person over the last 12 months?
[select one]
The purpose of this question is to help answer other questions such as whether treatment, survival and rehabilitation may differ according to employment status
(Insert country specific employment status list)

Information on acute stroke event

- | | | |
|--------|---|--|
| (I 21) | Date of stroke <i>Enter date of stroke symptoms onset, or first noticed</i> | [] [] / [] [] [] [] [] []
d d m m y y y y |
| (I 22) | Definite stroke?
<i>[select one]</i> | Yes (1) []
No (2)
Insufficient data (3) |
| (I 23) | If the patient has a definite stroke,
has the patient had a previous stroke?
<i>[select one]</i>
<i>Ensure accurate stroke diagnosis for previous stroke events</i> | Yes, original records seen (1) []
Yes, recorded in notes' history (2)
No (3)
Insufficient data (4) |
| (I 24) | If the patient has a definite stroke,
is this the first stroke in the surveillance period?
<i>[select one]</i> | Yes (1) []
No (2)
Insufficient data (3) |
| (I 25) | If No, was the previous stroke more than 28
days ago?
<i>[select one]</i>
Give date of the most recent previous stroke
since the surveillance period began.
<i>Enter date of stroke symptoms onset, or first noticed</i> | Yes (1) []
No (2)
Insufficient data (3)
[] [] / [] [] [] [] [] []
d d m m y y y y |

OPTIONAL items *(to be defined by centres; see comments section 3-9)*

(01) XX [_____]

(02) XX [_____]

Stroke Events Admitted To Hospital (Step 1)

For further guidance on Step 1, Events Admitted to Hospital; see Section 5, page 5-17

Hospital admission

(S1 1) Date of admission to hospital [][]/[][]/[][][][]
If in-hospital stroke, insert day of hospitalization from primary disease d d m m y y y y

(S1 2)	Which department(s)/ unit(s) was the patient treated in?	Intensive care unit	[]
	<i>[insert 1 for YES, 0 for NO, or 9 for UNKNOWN]</i>	Medical unit	[]
	<i>If the patient was treated in several units, list all. If Unknown, enter 9</i>	Neurological unit	[]
		Neurosurgical unit	[]
		Rehabilitation unit	[]
		Stroke unit	[]
		Other	[]

(S1 3)	What was the living situation of the patient pre stroke?	Independent at home	(1) []
	<i>[select one]</i>	Dependent at home	(2)
	<i>If in-hospital stroke patient, insert living situation prior to hospitalisation</i>	Community facility	(3)

(S1 4)	Modified Rankin scale prior to stroke <i>[select one]</i> <i>See Section 5-16 of the Stroke manual for further details on the Modified Rankin Scale.</i>	<i>No symptoms at all</i> <i>No significant disability</i> <i>despite symptoms</i> <i>Slight disability</i> <i>Moderate disability, but able</i> <i>to walk without assistance</i> <i>Moderate disability, but unable</i> <i>to walk without assistance</i> <i>Severe disability</i> <i>Unknown</i>	(0) [] (1) (2) (3) (4) (5) (9)
--------	--	--	--

(S1 5) Which of following neurological signs were present at first medical examination after hospitalization?
[insert 1 for YES, 0 for NO, or 9 for UNKNOWN]
Insert 1 if neurological sign was present at first medical examination
0 if sign was not present and 9 if Unknown.

Disturbed consciousness	[]
Weakness/ paresis	[]
Speech disturbances	[]

Stroke classification

(S1 6) What subtype of stroke was diagnosed? Ischemic stroke (1) []
[select one] Intracerebral hemorrhage (2)
See Section 1-5 for explanation of different types of stroke. Subarachnoid hemorrhage (3)
Unspecified type (4)

In-hospital assessment

- | | | |
|--|--------------------------------|-----|
| (S1 13) Which of the following assessments were done during the patients stay in hospital? | Seen by occupational therapist | [] |
| <i>[insert 1 for YES, 0 for NO, or 9 for UNKNOWN]</i> | Seen by physiotherapist | [] |
| | Seen by speech therapist | [] |
| | Swallowing assessment | [] |

Refers to assessments during hospitalization irrespective of whether the Patient was treated or not after the first visit.

Complications during hospitalization

- | | | |
|---|------------------------|-----|
| (S1 14) Which of the following complications occurred during the patients stay in hospital? | Deep venous thrombosis | [] |
| <i>[insert 1 for YES, 0 for NO, or 9 for UNKNOWN]</i> | Other CV complication | [] |
| | Pneumonia | [] |

Discharge from hospital

- (S1 15) What was the vital status at discharge? Patient alive (1) []
[select one] Patient dead (2)

- (S1 16) If patient died in hospital, indicate day of death [] [] / [] [] / [] [] [] []
This date is required to calculate early survival rates. d d m m y y y y

- (S1 17) If patient alive at discharge, indicate date of discharge [] [] / [] [] / [] [] [] []
This date is required to calculate survival rates and length of hospital stay. d d m m y y y y

- (S1 18) If patient alive at discharge, what was the discharge destination of the patient?
[select one]
- See Section 5-19 for further details
- | | |
|----------------------------|---------|
| Home | (1) [] |
| Other hospital/ other ward | (2) |
| Community facility | (3) |
| Unknown | (9) |

- | | | | |
|---------|--|--|---------|
| (S1 19) | If patient alive at discharge, | <i>No symptoms at all</i> | (0) [] |
| | Modified Rankin scale at discharge | <i>No significant disability</i> | |
| | <i>[select one]</i> | despite symptoms | (1) |
| | <i>See Section 5-16 for further details.</i> | <i>Slight disability</i> | (2) |
| | | <i>Moderate disability, but able</i> | |
| | | to walk without assistance | (3) |
| | | <i>Moderate disability, but unable</i> | |
| | | to walk without assistance | (4) |
| | | <i>Severe disability</i> | (5) |
| | | <i>Unknown</i> | (9) |

Follow up at 28 day after stroke onset (Optional)

- | | | |
|-------|--|--|
| (F 1) | Was it possible to follow up the patient at day 28?
[select one]
See Section 5-20 for further details | Yes (1) []
No, no contact (2)
No, patient refused (3) |
| (F 2) | If patient was followed up, indicate date of follow up | [] [] / [] [] / [] [] [] []
d d m m y y y y |
| (F 3) | How was the 28d follow up of the patient performed?
[select one]
See Section 5-20 for further details
Use "does not apply" for patients that died within the first 28 days,
refused participation or could not be contacted. | Medical records only (1) []
Physical examination (2)
Telephone interview (3)
Postal follow up (4)
Other (5)
Does not apply (6) |
| (F 4) | What is the vital status at day 28?
[select one]
Use "unknown" if no follow-up was performed | Patient alive (1) []
Patient dead (2)
Unknown (9) |
| (F 5) | If patient dead at day 28 indicate day of death | [] [] / [] [] / [] [] [] []
d d m m y y y y |
| (F 6) | If the patient alive at day 28, what is
the living situation of the patient at day 28?
[select one] | Home (1) []
Community facility (2)
Still in hospital (3) |
| (F 7) | If patient alive at day 28,
Modified Rankin scale at day 28
[select one] | No symptoms at all (0) []
No significant disability
despite symptoms (1)
Slight disability (2)
Moderate disability, but able
to walk without assistance (3)
Moderate disability, but unable
to walk without assistance (4)
Severe disability (5)
Unknown (9) |

Fatal Stroke Events in the Community (Step 2)

For further guidance on Step 2, Fatal Events in the Community; see Section 5-21

(S2 1) Indicate date of death

$$\begin{array}{cccccccc} [&] & [&] & / & [&] & [&] & / & [&] & [&] & [&] & [&] \\ \text{d} & \text{d} & & \text{m} & \text{m} & & \text{y} & \text{y} & \text{y} & \text{y} & & & & & & & & \end{array}$$

(S2 2) How was the patient managed in community from stroke onset until death?
[select one]

In nursing home	(1) []
At home by doctor	(2)
Other medical consultation	(3)
Medically unattended	(4)
Insufficient data	(5)

(S2 3) How was the information about fatal stroke events in the community collected?
[select one]

Verbal autopsy	(1) []
Death certificate	(2)
Medical autopsy	(3)

See Section 5-5 or further details these methods

(S2 4) If information was derived from **death certificate** which International Disease Classification (ICD) System was used?
[select one]

ICD 8 System	(1) []
ICD 9 System	(2)
ICD 10 System	(3)
No ICD System	(4)

See Section 5-21 for further details on the ICD system

(S2 5) If ICD System was used, indicate ICD code
See Section 5-21 for further details on ICD codes

See Section 5-21 for further details on ICD codes

(S2 6) If a **medical autopsy** was performed, what subtype of stroke was diagnosed?
[select one]

Ischemic stroke	(1) []
Intracerebral hemorrhage	(2)
Subarachnoid hemorrhage	(3)
Unspecified type	(4)

Non-Fatal Stroke Events in the Community (Step 3)

For further guidance on Step 3, Non-Fatal Events in the Community; see Section 5-7

- | | | | |
|--------|--|--|--|
| (S3 1) | How was the patient managed in community?
<i>[select one]</i> | In nursing home
Medically unattended
At home by doctor or nurse
Other medical consultation
Insufficient data | (1) []
(2)
(3)
(4)
(5) |
| (S3 2) | How was the information about the non-fatal stroke event in the community collected?
<i>[select one]</i> | Survey of health facilities
Survey of hemiplegia | (1) []
(2) |
| (S3 3) | What subtype of stroke was diagnosed?
<i>[select one]</i>
<i>See Section 1- 5 for further information on stroke subtypes</i> | Ischemic stroke
Intracerebral hemorrhage
Subarachnoid hemorrhage
Unspecified type | (1) []
(2)
(3)
(4) |
| (S3 4) | How was the diagnosis of stroke subtype verified?
<i>[select one]</i> | Clinical diagnosis alone
By diagnostic techniques | (1) []
(2) |
| (S3 5) | What was the living situation of the patient pre stroke?
<i>[select one]</i> | Independent at home
Dependent at home
Community facility
Unknown | (1) []
(2)
(3)
(9) |
| (S3 6) | Did the patient receive one or more of the following medications
<i>[insert 1 for YES, 0 for NO, or 9 for UNKNOWN]</i>
<i>See Section 5-19 for further details on medical treatment.</i> | Anticoagulant drugs
Antidiabetic drugs
Antiplatelet drugs
Cholesterol lowering drugs
Tablets for high blood pressure
Others | []
[]
[]
[]
[]
[] |



Registration of Sites participating in STEPS Stroke Surveillance

Introduction

This form registers an expression of interest in participating in STEPS Stroke and provides the ICS with background information on the planned stroke surveillance site.

Name of SSS

Site coordinator	Contact details
Title	
Name (family)	
Name (first)	
Institution	
Mail address	
City, Country	
Postal code	
Telephone	
Fax	
E-mail	

Other details

Please provide brief details of your experience in cerebrovascular disease and other details relating to data management:

Type	Previous experience (please tick or give brief details)
Clinical	
Research	
Epidemiologic	
Other	

Data management	Yes	No
Are computers available for data entry?		
Do you have software license for MicroSoft Access?		
Which version of MicroSoft Assess do you have?		
Is secure storage of original and electronic data possible?		
Do you take responsibility for all data entered?		
Do you have access to a data analyst?		

Ethical and legal issues	Yes	No
Does the site coordinator take responsibility for all ethical and legal issues related to the register (including approval from local ethics committee, insurance for staff members, providing informed consent, data security etc)?		
Financial support		
Have you obtained financial support to cover all expenses Related to the register? Please state the kind of financial support (in kind, Grant etc):		

Access to routine DCs Do you have access to routine death certificates?
Please give details on how you plan to verify potential stroke deaths.

Register duration Please provide details of proposed duration

Register duration	Please tick
Start date of Registration	
Duration - ongoing register	
Planned Start date (first enrolled patient)	

Proposed design Indicate which of the following register designs is planned:

Register design	Please tick	Estimated # of stroke patients per year
Hospital register - case series only		
Hospital register - with population denominator		
Population based incidence register		

Note: for meaningful analyses by age and sex, a minimum of 250 per year is required

Case finding Please provide details of planned case finding methods (including frequency)

Hot pursuit	
Cold pursuit	
Mixed pursuit	
Frequency	

Hospital registers

Please list all health facilities to be included in the proposed Stroke Surveillance Site

Hospital Name	Type	No. Wards (No. beds)	Est # stroke cases per week	Brain imaging available? (please tick)

Population based registers

If data are available from the source population from which all eligible stroke cases (Hospital register - step 1; or Incidence register - Steps 1-3), please complete:

Age group (yrs)	Instrument coverage	Men	Women	Total
0 - 14	Optional			
15 - 24	Expanded			
25 - 34	Expanded			
35 - 44	Core			
45 - 54	Core			
55 - 64	Core			
65 - 74	Core			
75 - 84	Core			
85 - 94	Expanded			
95 or more	Optional			
Total				

Population

Which of the following provides the information for your defined population?

Source	Date of info
Census	
Intercensal estimate	
Household registers	
Other population based register (eg cancer)	
Other (briefly explain)	

Final STEPS Instrument

Please send a copy of the planned instrument to be used in your site. This should include expanded items (and any optional items if you require feedback on format).

Signature

Date:

Section 8: Glossary and Reference Material

Introduction This section provides an alphabetical list of all the terms used in a STEPS surveillance with definitions that are appropriate for STEPS and a list of all the source and reference material used to compile this manual.

In this section This section covers the following topics.

Topic	See Page
Glossary of Terms used in STEPS Stroke	8-2
Source Publications and References	8-3

Glossary of Terms used in STEPS Stroke

Term	Definition
Amaurosis fugax	Periodical blindness of an eye due to embolic occlusion of the artery supplying the retina.
Apraxia	The inability to execute a planned motor act in the absence of paralysis of the muscles normally used in the performance of the act.
Ataxia	Co-ordination disturbances.
Bilateral	Includes both sides of the body.
Case-fatality	The proportion of events which are fatal within a given period of time.
Contra-lateral	Refers to the opposite side of the body.
Demography	The composition of the population.
Diplopia	Double vision.
Dysarthria	A defect in the articulation of the speech.
Dysphagia	Impaired ability to swallow.
Dysphasia	Difficulty with comprehension or production of the language despite intact articulation and phonation.
Hemiplegia	Weakness of the arm and leg on one side of the body.
Homonymous hemianopia	Loss of vision in one half of the visual field. Lesions of the optic nerve behind the chiasm produce contra-lateral visual field deficits.
Incidence	The incidence of stroke is the number of new cases of stroke arising in a given period in a defined population: usually expressed as a rate per 100,000. Incidence gives an indication of the risk of stroke.
Intracerebral hemorrhage	Bleeding from intracerebral arteries and may cause stroke symptoms.
Ischemic stroke	Stroke symptoms known to originate from an occlusion of cerebral arteries.
Modified Rankin Scale	A scale that indicates the level of handicap in a person.
Morbidity	A rate of how many people get sick per person years.
Mortality	A rate of how many people die per person years.
Prevalence	Prevalence is the number of cases in a defined population at a specified point in time - gives a "snapshot" of survivors at any one time
Stroke	A clinical diagnosis based on recognisable clinical symptoms indicating a vascular cause of sudden onset of neurological deficits. For definition please see page 1-4.
Subarachnoid hemorrhage	A bleeding from intra cranial arteries leading to blood between two membranes that surround the brain.
Surveillance	Ongoing, continuous collection of epidemiologic data in a population.
Transient Ischemic Attack (TIA)	Sudden neurologic deficits that lasts less than 24 hours, and with full recovery.
Unilateral	Restricted to one side of the body.
Vertigo	A false sense of rotary movement of self or surrounding objects. May be associated with nausea and vomiting.

Source Publications and References

Introduction

This section provides an alphabetical list of :

- References used in this publication
 - Resources available from the STEPS team
-

References

This section provides an alphabetical list of references and sources used.

Anderson C, Carter, K, Hackett M, Feigen V, Barber, A, Broad J, Bonita R. Trends in stroke incidence in Auckland, New Zealand, between 1981 and 2003. *Stroke* 2005; 36:2087.

Armstrong T, Bonita R. Building capacity for an integrated noncommunicable disease risk factor surveillance system in developing countries. *Ethnicity and Disease* 2003;13(s)2

Asplund K, Bonita, Kuulasmaa, Rajakangas A-M, Feigin V, Schaedlich H, Suzuki K, Thorvaldsen P, Tuomilehto J; for the WHO MONICA project. Multinational comparisons of stroke epidemiology: evaluation of case ascertainment in the WHO MONICA Stroke Study. *Stroke* 1995;26:355-60

Asplund, K., Rajakangas, A., Kuulasmaa, K., Thorvaldsen, P., Bonita, R., Stegmayr, B., Suzuki, K., and Eisenblätter, D. WHO MONICA Project: Multinational comparison of diagnostic procedures and management of acute stroke. *Cerebrovasc Dis* 1996; 6: 66-74.

Adams HP Jr, Adams R, Adams RJ, Brott T, Brott TF, del Zoppo GJ, Furlan et al. Guidelines for the early management of patients with ischemic stroke: A scientific statement from the Stroke Council of the American Stroke Association. *Stroke* 2003; 34:1056-83.

Bonita R, Mendis S, Truelsen T, Bogousslavsky J, Toole J, Yatsu F. The Global Stroke Initiative. *Lancet Neurology*. 2004;3:391-93

Bonita R, Truelsen T. Stroke in Sub Saharan Africa: a neglected chronic disease. *Lancet Neurology*. 2003; 2: 592-4.

Bonita, R; Douglas, K; Winkelmann, R; De Courten, M; The WHO STEPwise approach to surveillance (STEPS) of noncommunicable disease risk factors. Chapter in (eds) McQueen, DV and Puska, P (editors); *Global Risk Factor Surveillance*. London: Kluwer Academic/Plenum Publishers, New York. 2003; pp 9-22

Continued on next page

Source Publications and References, Continued

Bonita, R; Truelsen, T. Edited by Barnett, H.J.M; Bogousslavsky, J, Meldrum, H. Advances in Ischemic Stroke Epidemiology. Chapter in Ischemic Stroke: Advances in Neurology. Vol 92. Lippincott Williams & Wilkins, Philadelphia 2003

Bonita, R; Strong, K; De Courten, M.; From surveys to surveillance. Rev Panam Salud Publica/Pan Am J Public Health; 2001; 10: 223–225

Bonita R, Broad JB, Anderson NE, and Beaglehole R. Approaches to the problems of measuring the incidence of stroke. Int J Epidemiol 1995;24: 535-542.

Bharucha, NE, Bharucha, EP, Bharucha AE, Bhise AV, Schoenberg BS.. Prevalence of stroke in the parsi community of Bombay. Stroke 1998; 19: 60-62.

Choi B, Corber, McQueen D, Bonita R et al. Enhancing regional capacity in chronic disease surveillance in the Americas. Rev Panam Salud Publica/ Pan Am J Public Health 2005;17: 130-141.

Prospective studies collaboration. Cholesterol, diastolic blood pressure, and stroke: 13,000 strokes in 450,000 people in 45 prospective cohorts. - Lancet 1995; 346:1647-53.

Collins RF, Armitage JF, Parish S, Sleight P, Sleight PF, Peto R. - Effects of cholesterol-lowering with simvastatin on stroke and other major vascular events in 20536 people with cerebrovascular disease or other high-risk conditions. - Lancet 2004; 363:757-67.

Chandramohan D, Maude GH, Rodrigues LC, and Hayes RJ. Verbal autopsies for adult deaths: issues in their development and validation. Int J Epidemiol 1994; 23: 213-222.

Dennis MS, Bamford JM, Molyneux AJ, and Warlow CP. Rapid resolution of signs of primary intracerebral haemorrhage in computed tomograms of the brain. BMJ 1987; 295:379-381.

Dhamija RK, Mittal S, and Bansal BC. Trends in clinico-epidemiological correlates of stroke in the community. J Indian Academy Clin Med. 2000;5:27-31

Doggett DL, Tappe KA, Mitchell MD, Chapell RF, Coates VF, Turkelson CM. Prevention of pneumonia in elderly stroke patients by systematic diagnosis and treatment of dysphagia: an evidence-based comprehensive analysis of the literature. Dysphagia 2001; 16:279-95.

Continued on next page

Source Publications and References, Continued

D'Olhaberriague LF, Litvan IF, Mitsias PF, Mansbach HH. A reappraisal of reliability and validity studies in stroke. *Stroke* 1996; 27:2331-6.

Donnan GA - Davis S, Davis SM. Stroke and cholesterol: weakness of risk versus strength of therapy. *Stroke* 2004; 35:1526.

Feng J H, Nowson CA, MacGregor GA. Fruit and vegetable consumption and stroke: a met analyses of cohort studies. *Lancet* 2006; 367:320-326

Feigin VL, Carter K. Editorial comment. Stroke incidence studies one step closer to the elusive gold standard? *Stroke*. 2004; 35: 2045-7.

Feigin V, Brainin M, Breteler MMB, Marytn C, Wolfe C, Bornstein N. Fieschi C et al . Teaching in neuro epidemiology in Europe: Time to get serious. *Eur J Neurology*; 2004; 11:795-9

Feigin VL, Lawes CM, Bennett DA, Anderson CS. Stroke epidemiology: a review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurol*. 2003; 2: 43-53. Review.

Foerch C, Misselwitz B, Sitzler M, Berger K, Steinmetz H, Neumann-Haefelin T. Difference in recognition of right and left hemispheric stroke. *Lancet* 2005; 366: 392-393.

Hatano S. Experience from a multi centre stroke register: a preliminary report. *Bull WHO* 1975; 54:541-553.

Heuschmann PU, Kolominsky-Rabas P, Kolominsky-Rabas PL, Roether J, Roether JF, Misselwitz BF, Lowitzsch KF, et al. - Predictors of in-hospital mortality in patients with acute ischemic stroke treated with thrombolytic therapy. *JAMA* 2004; 292:1831-8.

Heuschmann PU, Kolominsky-Rabas P, - Kolominsky-Rabas PL FAU, Misselwitz B, Misselwitz BF, Hermanek PF, Leffmann CF, et al. - Predictors of in-hospital mortality and attributable risks of death after ischemic stroke: the German Stroke Registers Study Group. - *Arch Intern Med* 2004 13; 164:1761-8.

Katzan IL, Cebul RD FAU, Husak SH, Dawson NV, Baker DW. - The effect of pneumonia on mortality among patients hospitalized for acute stroke. - *Neurology* 2003; 60:620-5.

Lavados PM, Sacks C, Prina L, Escobar A, Tossi C, Araya F, Feuerhake W, Galvez M, Salinas R, Alvarez G. Incidence, 30 day case-fatality rates and prognosis of stroke in Iquique, Chile (PISCIS project). *Lancet* 2005;365,:2206-2215

Continued on next page

Source Publications and References, Continued

Law MR, Wald NJ, Rudnicka AR. Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis. - BMJ 2003; 326:1423.

Murray CJL and Lopez AD. Mortality by cause for eight regions of the world: Global Burden of Disease Study. Lancet. 1997; 349, 1269-1276.

Nicoletti A, Sofia V, Giuffrida S, Bartoloni A, Bartalesi F, Bartolo MLL, Fermo SL, Cocuzza V, Gamboa H, Salazar E, and Reggio A. Prevalence of stroke. A door-to-door survey in rural Bolivia. Stroke 2000;31: 882-885.

Nor AM, Davis J, Sen B, Shipsey D, Louw SJ, Dyker AG, Davis M, Ford GA. The Recognition of Stroke in the Emergency Room (ROSIER) scale: development and validation of a stroke recognition instrument. Lancet Neurology. 2005; 4:727-734.

Pandian JD, Jaison A, Sukhbinder SD, Kalra G, Shamsheer S, Lincoln DJ, Abraham G. Public awareness of warning symptoms, risk factors, and treatment of stroke in Northwest India. Stroke 2005; 36: 644-648.

Pendlebury ST, Rothwell PM, Algra A, Ariesen M-J, Bakac G, Czlonkowska A, et al. Underfunding of stroke research: a European wide problem. Stroke 2004; 35:2368-71

Poungvarin N. Stroke in the developing world. Lancet 1998; 35:19-22.

Piechowski-Jozwiak BF, Bogousslavsky J. Cholesterol as a risk factor for stroke: the fugitive? Stroke 2004;35:1523

Reed SD, Cramer SC, Blough DK, Meyer KF, Jarvik JG. - Treatment with tissue plasminogen activator and inpatient mortality rates for patients with ischemic stroke treated in community hospitals. Stroke 2001; 32:1832-40.

Rothwell P, Coull A, Silver L, Fairhead J, Giles M, Lovelock C, Redgrave J, et al. Population-based study of event-rate, incidence, case fatality, and mortality for all acute vascular events in all arterial territories (Oxford Vascular Study) Lancet. 2005; 366: 1773-1783.

Sacco RL, Benjamin EJ FAU, Broderick JP FAU, Dyken MF, Easton JD, Feinberg WM et al. - AHAPrevention Conference. IV. Prevention and Rehabilitation of Stroke. Risk factors. - Stroke 1997; 28:1507-17.

Setel PW, Sankoh O, Rao C, Velkoff VV, Mathers, Gonghuan Z, Hemed et al. Sample registration of vital events with verbal autopsy. A renewed commitment to measuring and monitoring vital statistics. Bulletin WHO 2005; 83: 611-17.

Continued on next page

Source Publications and References, Continued

Straus SE, Majumdar S, Majumdar SR, McAlister F, McAlister FA. New evidence for stroke prevention: clinical applications. *JAMA* 2002;288:1396-98.

Stroke Unit Trialists' Collaboration. Collaborative systematic review of the randomized trials of organized inpatient (stroke unit) care after stroke. *BMJ* 1997;314: 1151-1159.

Struijs JN, van Genugten ML, Evers SM, Ament AJ, Baan CA, van den Bos GA. Modelling the future burden of stroke in The Netherlands: impact of aging, smoking, and hypertension. *Stroke* 2005; 36:1648-55.

Sudlow CL, Warlow CP. Comparable studies of the incidence of stroke and its pathological types: results from an international collaboration. International Stroke Incidence Collaboration. *Stroke*. 1997; 28:491-9.

Thorvaldsen P, Asplund K, Kuulasmaa K, Rajakangas A-M, Schroll M; for the WHO MONICA Project. *Stroke*, 1995;26: 361-67.

Truelsen T, Bonita R. *Surveillance of Stroke: A Global Perspective*. *Int J Epidemiol* 2001; 30; Suppl 1: S11-S16

Truelsen T, Mähönen M, Tolonen H, Asplund K, Bonita R, Vanuzzo D, for the WHO MONICA Project: *Trends in Stroke and Coronary Heart Disease in the WHO MONICA Project*. *Stroke, JAMA*. 2003; 34: 1346-1352.

UK TIA Study Group. The United Kingdom transient ischaemic attack (UK TIA). Aspirin Trial: Final results. *J Neurol Neurosurg Psychiatry*. 1991; 54:1044-1054.

Walker RW, McLarty DG, Kitange HM, Whiting D, Masuki G, Mtasiwa DM, Machibya H, Unwin N, and Alberti KGMM. Stroke mortality in urban and rural Tanzania. *Lancet* 2000;355:1684

Weir NU, Sandercock PA FAU, Lewis SC FAU, Signorini DF, Warlow CP. - Variations between countries in outcome after stroke in the International Stroke Trial (IST). - *Stroke* 2001; 32:1370-7.

Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation: a major contributor to stroke in the elderly. The Framingham Study. *Arch Intern Med* 1987; 147:1561-4

Wolfe CD, Tilling KF, Beech RF, Rudd AG. - Variations in case fatality and dependency from stroke in western and central Europe. The European BIOMED Study of Stroke Care Group. - *Stroke* 1999; 30:350-6.
