Paraguay - Asuncion (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Paraguay - Asuncion GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Paraguay - Asuncion could include in a comprehensive tobacco control program.

The Paraguay - Asuncion GYTS was a school-based survey of students in forms 7, 8 and 9 conducted in 2008.

A two-stage cluster sample design was used to produce representative data for Paraguay - Asuncion. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96.0%, the class response rate was 100.0%, the student response rate was 88.4%, and the overall response rate was 84.9%. A total of 3,275 students aged 13-15 participated in the Paraguay - Asuncion GYTS.

Prevalence

25.0% of students had ever smoked cigarettes (Boy = 27.7%, Girl = 22.5%)

18.1% currently use any tobacco product (Boy = 20.2%, Girl = 16.0%)

10.4% currently smoke cigarettes (Boy = 11.7%, Girl = 9.3%)

10.1% currently use other tobacco products (Boy = 11.8%, Girl = 8.6%)

17.6% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

13.1% think boys and 8.0% think girls who smoke have more friends 9.4% think boys and 4.9% think girls who smoke look more attractive

Access and Availability - Current Smokers

13.7% usually smoke at home

43.0% buy cigarettes in a store

80.3% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

36.1% live in homes where others smoke in their presence

56.2% are around others who smoke in places outside their home

85.3% think smoking should be banned from public places

66.8% think smoke from others is harmful to them

32.5% have one or more parents who smoke

11.2% have most or all friends who smoke

Cessation - Current Smokers

59.4% want to stop smoking

75.0% tried to stop smoking during the past year

75.3% have ever received help to stop smoking

Media and Advertising

83.7% saw anti-smoking media messages, in the past 30 days

90.9% saw pro-cigarette ads on billboards, in the past 30 days

88.3% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

12.1% have an object with a cigarette brand logo

15.4% were offered free cigarettes by a tobacco company representative

School

71.0% had been taught in class, during the past year, about the dangers of smoking 39.7% had discussed in class, during the past year, reasons why people their age smoke 62.8% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 18.1% of students currently use any form of tobacco; 10.4% currently smoke cigarettes; 10.1% currently use some other form of tobacco.
- SHS exposure is high over one-third of the students live in homes where others smoke, and close to 3 in 5 students are exposed to smoke around others outside of the home; more than 3 in 10 students have at least one parent who smokes.
- Two-thirds of the students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Six in 10 current smokers want to stop smoking.
- One in 8 students has an object with a cigarette brand logo on it.
- More than 4 in 5 students saw anti-smoking media messages in the past 30 days; 9 in 10 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.