Peru - Lima (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Peru - Lima GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Peru - Lima could include in a comprehensive tobacco control program.

The Peru - Lima GYTS was a school-based survey of students in 1^{st} , 2^{nd} , 3^{rd} , 4^{th} and 5^{th} secondary conducted in 2003.

A two-stage cluster sample design was used to produce representative data for Peru - Lima. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 96.0%, the class response rate was 100.0%, the student response rate was 90.3%, and the overall response rate was 86.7%. A total of 1,045 students aged 13-15 participated in the Peru - Lima GYTS.

Prevalence

59.6% of students had ever smoked cigarettes (Boy = 63.9%, Girl = 54.5%)

23.4% currently use any tobacco product (Boy = 24.4%, Girl = 21.6%)

19.2% currently smoke cigarettes (Boy = 20.8%, Girl = 17.0%)

7.9% currently use other tobacco products (Boy = 8.2%, Girl = 7.1%)

30.5% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

14.8% think boys and 15.1% think girls who smoke have more friends 11.9% think boys and 12.2% think girls who smoke look more attractive

Access and Availability - Current Smokers

10.0% usually smoke at home

59.3% buy cigarettes in a store

73.9% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

25.1% live in homes where others smoke in their presence

41.7% are around others who smoke in places outside their home

85.9% think smoking should be banned from public places

56.6% think smoke from others is harmful to them

41.1% have one or more parents who smoke

16.9% have most or all friends who smoke

Cessation - Current Smokers

62.2% want to stop smoking

64.5% tried to stop smoking during the past year

58.6% have ever received help to stop smoking

Media and Advertising

90.6% saw anti-smoking media messages, in the past 30 days

76.9% saw pro-cigarette ads on billboards, in the past 30 days

82.0% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

12.5% have an object with a cigarette brand logo

9.7% were offered free cigarettes by a tobacco company representative

School

43.2% had been taught in class, during the past year, about the dangers of smoking

37.1% had discussed in class, during the past year, reasons why people their age smoke

32.2% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 23.4% of students currently use any form of tobacco; 19.2% currently smoke cigarettes; 7.9% currently use some other form of tobacco.
- SHS exposure is high onequarter of the students live in homes where others smoke, and more than 2 in 5 students are exposed to smoke around others outside of the home; 2 in 5 students have at least one parent who smokes.
- Close to 3 in 5 students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- More than 6 in 10 current smokers want to stop smoking.
- One in 8 students has an object with a cigarette brand logo on it.
- Nine in 10 students saw antismoking media messages in the past 30 days; close to 4 in 5 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.