Peru - Tarapoto (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Peru - Tarapoto GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Peru - Tarapoto could include in a comprehensive tobacco control program.

The Peru - Tarapoto GYTS was a school-based survey of students in 1^{st} , 2^{nd} , 3^{rd} , 4^{th} and 5^{th} secondary conducted in 2003.

A two-stage cluster sample design was used to produce representative data for Peru - Tarapoto. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 100.0%, the student response rate was 92.1%, and the overall response rate was 92.1%. A total of 1,126 students aged 13-15 participated in the Peru - Tarapoto GYTS.

Prevalence

48.7% of students had ever smoked cigarettes (Boy = 56.1%, Girl = 40.8%)

18.8% currently use any tobacco product (Boy = 22.9%, Girl = 14.3%)

15.5% currently smoke cigarettes (Boy = 20.4%, Girl = 10.0%)

6.4% currently use other tobacco products (Boy = 6.7%, Girl = 6.1%)

19.6% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

21.1% think boys and 17.5% think girls who smoke have more friends 13.3% think boys and 11.5% think girls who smoke look more attractive

Access and Availability - Current Smokers

19.5% usually smoke at home

53.9% buy cigarettes in a store

77.2% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

27.2% live in homes where others smoke in their presence

35.5% are around others who smoke in places outside their home

88.7% think smoking should be banned from public places

58.1% think smoke from others is harmful to them

42.3% have one or more parents who smoke

10.3% have most or all friends who smoke

Cessation - Current Smokers

77.6% want to stop smoking

80.5% tried to stop smoking during the past year

75.2% have ever received help to stop smoking

Media and Advertising

91.8% saw anti-smoking media messages, in the past 30 days

74.4% saw pro-cigarette ads on billboards, in the past 30 days

78.4% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

10.6% have an object with a cigarette brand logo

8.9% were offered free cigarettes by a tobacco company representative

School

64.9% had been taught in class, during the past year, about the dangers of smoking 55.3% had discussed in class, during the past year, reasons why people their age smoke 50.4% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 18.8% of students currently use any form of tobacco; 15.5% currently smoke cigarettes; 6.4% currently use some other form of tobacco.
- SHS exposure is high more than one-quarter of the students live in homes where others smoke, and over one-third of the students are exposed to smoke around others outside of the home; more than 2 in 5 students have at least one parent who smokes.
- Close to 3 in 5 students think smoke from others is harmful to them.
- Almost 9 in 10 students think smoking in public places should be banned.
- Over three-quarters of the current smokers want to stop smoking.
- One in 10 students has an object with a cigarette brand logo on it.
- More than 9 in 10 students saw anti-smoking media messages in the past 30 days; over three-quarters of the students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.