Venezuela - Cojedes (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Venezuela - Cojedes GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Venezuela - Cojedes could include in a comprehensive tobacco control program.

The Venezuela - Cojedes GYTS was a school-based survey of students in grades 6, 7, 8 and 9 conducted in 2003.

A two-stage cluster sample design was used to produce representative data for Venezuela - Cojedes. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 100.0%, the student response rate was 78.8%, and the overall response rate was 78.8%. A total of 902 students aged 13-15 participated in the Venezuela - Cojedes GYTS.

Prevalence

18.6% of students had ever smoked cigarettes (Girl = 19.1%, Girl = 17.7%)

12.5% currently use any tobacco product (Girl = 14.0%, Girl = 11.2%)

6.5% currently smoke cigarettes (Girl = 4.8%, Girl = 7.9%)

7.4% currently use other tobacco products (Girl = 11.0%, Girl = 4.2%)

17.0% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

9.2% think boys and 8.0% think girls who smoke have more friends 6.7% think boys and 5.5% think girls who smoke look more attractive

Access and Availability - Current Smokers

22.3% usually smoke at home

34.3% buy cigarettes in a store

Exposure to Secondhand Smoke (SHS)

40.4% live in homes where others smoke in their presence

51.3% are around others who smoke in places outside their home

89.4% think smoking should be banned from public places

62.4% think smoke from others is harmful to them

37.9% have one or more parents who smoke

8.3% have most or all friends who smoke

Cessation - Current Smokers

61.9% tried to stop smoking during the past year 76.5% have ever received help to stop smoking

Media and Advertising

79.1% saw anti-smoking media messages, in the past 30 days

71.7% saw pro-cigarette ads on billboards, in the past 30 days

70.8% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

9.7% have an object with a cigarette brand logo

11.1% were offered free cigarettes by a tobacco company representative

School

47.4% had been taught in class, during the past year, about the dangers of smoking

33.3% had discussed in class, during the past year, reasons why people their age smoke

50.1% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 12.5% of students currently use any form of tobacco; 6.5% currently smoke cigarettes; 7.4% currently use some other form of tobacco.
- SHS exposure is high 4 in 10 students live in homes where others smoke, and half the students are exposed to smoke around others outside of the home; over one-third of the students have at least one parent who smokes.
- More than 6 in 10 students think smoke from others is harmful to them.
- Nine in 10 students think smoking in public places should be banned.
- One in 10 students has an object with a cigarette brand logo on it.
- Eight in 10 students saw antismoking media messages in the past 30 days; over 7 in 10 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.