Ecuador - Zamora (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Ecuador - Zamora GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Ecuador - Zamora could include in a comprehensive tobacco control program.

The Ecuador - Zamora GYTS was a school-based survey of students in grades 1, 2 and 3 conducted in 2001.

A two-stage cluster sample design was used to produce representative data for Ecuador - Zamora. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 100.0%, the student response rate was 89.9%, and the overall response rate was 89.9%. A total of 716 students aged 13-15 participated in the Ecuador - Zamora GYTS.

Prevalence

59.4% of students had ever smoked cigarettes (Boy = 65.1%, Girl = 53.5%)

32.9% currently use any tobacco product (Boy = 32.9%, Girl = 30.8%)

24.8% currently smoke cigarettes (Boy = 27.4%, Girl = 22.0%)

17.6% currently use other tobacco products (Boy = 18.9%, Girl = 16.5%)

23.1% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

26.2% think boys and 19.8% think girls who smoke have more friends 13.5% think boys and 11.8% think girls who smoke look more attractive

Access and Availability - Current Smokers

21.8% usually smoke at home

46.4% buy cigarettes in a store

80.5% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

41.8% live in homes where others smoke

61.7% are around others who smoke in places outside their home

81.5% think smoking should be banned from public places

64.9% think smoke from others is harmful to them

46.3% have one or more parents who smoke

19.3% have most or all friends who smoke

Cessation - Current Smokers

80.1% want to stop smoking

64.9% tried to stop smoking during the past year

71.8% have ever received help to stop smoking

Media and Advertising

74.3% saw anti-smoking media messages, in the past 30 days

69.5% saw pro-cigarette ads on billboards, in the past 30 days

68.8% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

10.9% have an object with a cigarette brand logo

12.8% were offered free cigarettes by a tobacco company representative

School

54.9% had been taught in class, during the past year, about the dangers of smoking

22.5% had discussed in class, during the past year, reasons why people their age smoke

44.5% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 32.9% of students currently use any form of tobacco; 24.8% currently smoke cigarettes; 17.6% currently use some other form of tobacco.
- SHS exposure is high over 2 in 5 students live in homes where others smoke, and more than 3 in 5 students are exposed to smoke around others outside of the home; close to half the students have at least one parent who smokes.
- Close to two-thirds of the students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Four in 5 current smokers want to stop smoking.
- One in 9 students has an object with a cigarette brand logo on it
- Three-quarters of the students saw anti-smoking media messages in the past 30 days; 7 in 10 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.