Venezuela - Zulia (Ages 13-15) Global Youth Tobacco Survey (GYTS)



The Venezuela - Zulia GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Venezuela - Zulia could include in a comprehensive tobacco control program.

The Venezuela - Zulia GYTS was a school-based survey of students in grades 6, 7, 8 and 9 conducted in 2001.

A two-stage cluster sample design was used to produce representative data for Venezuela - Zulia. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 73.1%, the class response rate was 100.0%, the student response rate was 76.9%, and the overall response rate was 56.2%. A total of 825 students aged 13-15 participated in the Venezuela - Zulia GYTS.

Prevalence

- 22.7% of students had ever smoked cigarettes (Male = 26.9%, Female = 19.7%)
- 23.5% currently use any tobacco product (Male = 21.1%, Female = 23.9%)
- 9.6% currently smoke cigarettes (Male = 13.5%, Female = 6.4%)
- 16.1% currently use other tobacco products (Male = 11.2%, Female = 18.5%)
- 16.5% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

15.1% think boys and 9.7% think girls who smoke have more friends 8.2% think boys and 7.6% think girls who smoke look more attractive

Access and Availability - Current Smokers

19.6% usually smoke at home

50.6% buy cigarettes in a store

80.6% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

43.8% live in homes where others smoke in their presence

45.0% are around others who smoke in places outside their home

85.0% think smoking should be banned from public places

56.8% think smoke from others is harmful to them

40.7% have one or more parents who smoke

12.5% have most or all friends who smoke

Cessation - Current Smokers

70.2% want to stop smoking

67.2% tried to stop smoking during the past year

64.4% have ever received help to stop smoking

Media and Advertising

81.7% saw anti-smoking media messages, in the past 30 days

79.9% saw pro-cigarette ads on billboards, in the past 30 days

81.1% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

18.1% have an object with a cigarette brand logo

17.6% were offered free cigarettes by a tobacco company representative

School

42.3% had been taught in class, during the past year, about the dangers of smoking

32.6% had discussed in class, during the past year, reasons why people their age smoke

41.4% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 23.5% of students currently use any form of tobacco; 9.6% currently smoke cigarettes; 16.1% currently use some other form of tobacco.
- SHS exposure is high Over 2 in 5 students live in homes where others smoke, and close to half of the students are exposed to smoke around others outside of the home; 4 in 10 students have at least one parent who smokes.
- More than half the students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Seven in 10 current smokers want to stop smoking.
- More than 1 in 5 students have an object with a cigarette brand logo on it.
- Over 8 in 10 students saw antismoking media messages in the past 30 days; 4 in 5 students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.