



Epidemiological Alert:

New Confirmed Cases of Saint Louis Encephalitis in Capital City and Province of Buenos Aires, Risk of Viral Circulation

(31 March 2010)

The Ministry of Health of Argentina had confirmed new cases of Saint Louis Encephalitis (SLE) virus infection. Until 29 March 2010, it has been reported four confirmed cases of SLE infection in the nation's capital city and in Buenos Aires province. The SLE virus had previously caused outbreaks in the central part of the country (mainly in the provinces of Córdoba and Entre Ríos), and this is the first time that it has been detected in Buenos Aires. This virus has epidemic potential, as has been shown by what occurred in 1975 in the United States in the Ohio and Mississippi River basins, where some 2,000 people developed the neuro-invasive form of the disease.

Faced with this situation and with the verification of cases having occurred in Argentina, it has become necessary to strengthen surveillance of febrile syndrome as well as febrile syndrome with neurological manifestations, in order to provide guidance on the vector control strategy.

Current Situation

From 1 March 2010 to date, the Ministry of Health of Argentina has reported a total of 43 cases throughout the country:

- 4 of them confirmed,
- 9 probable,
- 17 suspected cases under investigation, and
- 13 discarded.

Of the confirmed cases, one involves a resident of the city of Buenos Aires and three involve residents of the province of Buenos Aires.

Saint Louis Encephalitis

The virus is part of a group of acute viral diseases of limited duration that affect the central nervous system (CNS).

Infectious Agent: The SLE virus belongs to the family *Flaviviridae*, genus *Flavivirus*.

Transmission: The SLE virus presents a mosquito-bird-mosquito cycle involving domestic and wild birds and mosquitoes of the *Culex sp.* species. Humans and domestic mammals can acquire the infection through the bite of an infected mosquito, but they do not act effectively as hosts to perpetuate transmission.

Epidemiology and Geographical Distribution: All inhabitants of and visitors to areas with SLE circulation are at risk of contracting the disease. The virus is distributed throughout the American continent, with the last occurrence being reported in Argentina in 2005–2006.

Incubation Period: Varies from 5 to 15 days.

Symptomology: Less than 1% of SLE cases develop clinical manifestations. Severe infections present with an abrupt onset of headache, high fever, dizziness, nausea, and malaise. The majority of cases recover spontaneously; however, some develop signs of infection of the CNS, including stiff neck, confusion, disorientation, tremors, and stupor, progressing even to coma. Almost 90% of older adults who acquire the disease develop encephalitis. Case fatality proportion ranges from 5 to 15%, increasing with age.

Treatment: There is no vaccine or specific antiviral treatment. In those cases that present clinical manifestations, appropriate supportive therapy should be administered.

Laboratory Diagnosis: Diagnostic methods include serology, viral isolation, and PCR, which is only carried out in specialized laboratories.

Prevention: Preventive measures focus on individual protection by applying mosquito repellent, destroying breeding sites, and protecting living spaces with mosquito netting. In areas with a high density of *Culex sp.*, chemical vector control should be considered.

Confirmed, Probable, and Suspected Cases of Saint Louis Encephalitis by province and place of residence. Argentina, 1 January –29 March 2010.

Province	Confirmed Cases	Probable Cases	Suspected Cases under Investigation
Buenos Aires (Bs As), greater Metropolitan area	3		
Ciudad de Bs As, city proper	1	5	1
Entre Rios		1	6
Santa Fe		2	2
Chaco			2
Catamarca			1
Cordoba		1	4
Corrientes			1
Total	4	9	17

Source: Report from the National Focal Point of Argentina, 29 March 2010.

According to the available information about all the cases in the city and in the province of Buenos Aires, 13 cases were reported as suspect for dengue febrile syndrome and 6 cases have presented neurological involvement (meningoencephalitis).

The average age of these cases is 34 years (range: 12–72 years), and 79% of the cases occurred among males.

Regarding the geographical location of the cases in the city of Buenos Aires, the confirmed and probable cases are located in five communes in the southern and southeastern parts of the city. The cases of the province of Buenos Aires were located in the southern and northern parts.

Investigation, prevention and control activities undertaken by the country include the following: vector control activities with isolation of suspected cases, taking samples from persons living with anyone reported as being a case, field case-finding of febrile cases by community agents and technical personnel, and issuing epidemiological alerts to intensity surveillance.

Recommendations

a. Intensify epidemiological surveillance and investigation activities

Strengthen surveillance of febrile syndrome and febrile syndrome with neurological manifestations, and conduct epidemiological investigation of detected cases in order to provide guidance on what strategies to follow for vector control.

b. Take preventive and control measures and evaluate impact

Ensure widespread use of personal protection to reduce the risk of exposure to the vector and take the necessary vector control measures.

Bibliographical References

1. National Alert and Rapid Response Center, Epidemiological Directorate of the Ministry of Health of Argentina, event verification report, SLI, N5 EW 11 2010 (Centro Nacional Alerta y Respuesta Rápida, Dirección de Epidemiología, Ministerio de Salud de Argentina. Encefalitis de San Luis. Informe de verificación de eventos N 5 SE 11 2010).
2. Centers for Disease Control and Prevention. Saint Louis Encephalitis Virus: Technical Information. Available at: <http://www.cdc.gov/sle/index.html>
3. Epidemiological Directorate of the Ministry of Health and the Environment of Argentina (Dirección de Epidemiología, Ministerio de Salud y Ambiente de Argentina.) Virus de la Encefalitis de San Luis en Córdoba. Riesgo en zonas templadas y subtropicales con abundancia del vector. Direpi\Alerta\A83.3\5.2005. [Saint Louis Encephalitis Virus in Córdoba. Vector Risk in Temperate and Subtropical Zones].
4. American Public Health Association. Control of Communicable Diseases Manual. 18th ed. Washington D.C. 2004.
5. Epidemiological Directorate of the Ministry of Health of Argentina (Dirección de Epidemiología, Ministerio de Salud de Argentina) Virus de la Encefalitis de St. Louis (SLE) CABA-Buenos Aires-Córdoba. Direpi\Alerta\A83.3\2010. [Saint Louis Encephalitis Virus in CABA-Buenos Aires-Córdoba].